arXiv:math/0501474v1 [math.AT] 26 Jan 2005

HOMOTOPY FIXED POINTS FOR $L_{K(n)}(E_n \wedge X)$ USING THE CONTINUOUS ACTION

DANIEL G. DAVIS*

Abstract. Let G be a closed subgroup of G_n, the extended Morava stabilizer group. Let E_n be the Lubin-Tate spectrum, X an arbitrary spectrum with trivial G-action, and let $\hat{L} = L_{K(n)}$. We prove that $\hat{L}(E_n \wedge X)$ is a continuous G-spectrum with homotopy fixed point spectrum $(\hat{L}(E_n \wedge X))^{hG}$, defined with respect to the continuous action. Also, we construct a descent spectral sequence whose abutment is $\pi_*((\hat{L}(E_n \wedge X))^{hG})$. We show that the homotopy fixed points of $\hat{L}(E_n \wedge X)$ come from the $K(n)$-localization of the homotopy fixed points of the spectrum $(F_n \wedge X)$.

1. Introduction

Let E_n be the Lubin-Tate spectrum with $E_{n*} = W[\![u_1, ..., u_{n-1}]\!][u^{\pm 1}]$, where the degree of u is -2, and the complete power series ring over the Witt vectors $W = W(\mathbb{F}_{p^n})$ is in degree zero. Let S_n denote the nth Morava stabilizer group, the automorphism group of the Honda formal group law Γ_n of height n over $\mathbb{F}_{p^n}$. Then let $G_n = S_n \rtimes \mathrm{Gal}$, where Gal is the Galois group $\mathrm{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$, and let G be a closed subgroup of G_n. Morava's change of rings theorem yields a spectral sequence

$$H^*_c(G_n; \pi_*(E_n \wedge X)) \Rightarrow \pi_* \hat{L}(X), \tag{1.1}$$

where the E_2-term is continuous cohomology, X is a finite spectrum, and $\hat{L}$ is Bousfield localization with respect to $K(n)_* = \mathbb{F}_p[v_n, v_n^{-1}]$, where $K(n)$ is Morava K-theory (see [15, Prop. 7.4], [32]). Using the G_n-action on E_n by maps of commutative S-algebras (work of Goerss and Hopkins ([12], [13]), and Hopkins and Miller [34]), Devinatz and Hopkins [6] construct spectra $E_n^{\mathrm{h}G}$ with strongly convergent spectral sequences

$$H^s_c(G; \pi_t(E_n \wedge X)) \Rightarrow \pi_{t-s}(E_n^{\mathrm{h}G} \wedge X). \tag{1.2}$$

Also, they show that $E_n^{\mathrm{h}G_n} \simeq \hat{L}(S^0)$, so that $E_n^{\mathrm{h}G_n} \wedge X \simeq \hat{L}(X)$.

We recall from [6, Rk. 1.3] the definition of the continuous cohomology that appears above (see also Lemma 2.16). Let $I_n = (p, u_1, ..., u_{n-1}) \subset E_{n*}$. The isomorphism $\pi_t(E_n \wedge X) = \lim_k \pi_t(E_n \wedge X)/I_n^k \pi_t(E_n \wedge X)$ presents $\pi_t(E_n \wedge X)$ as the inverse limit of finite discrete G-modules. Then the above continuous cohomology is defined by

$$H^s_c(G; \pi_t(E_n \wedge X)) = \lim_k H^s_c(G; \pi_t(E_n \wedge X)/I_n^k \pi_t(E_n \wedge X)).$$

Date: January 12, 2005.

*The author was supported by NSF grant DMS-9983601.

We compare the spectrum $E_n^{\mathbf{h}G}$ and spectral sequence (1.2) with constructions for homotopy fixed point spectra. When K is a discrete group and Y is a K-spectrum of topological spaces, there is a homotopy fixed point spectrum $Y^{hK} = \mathrm{Map}_K(EK_+, Y)$, where EK is a free contractible K-space. Also, there is a descent spectral sequence

$$E_2^{s,t} = H^s(K; \pi_t(Y)) \Rightarrow \pi_{t-s}(Y^{hK}),$$

where the E_2-term is group cohomology [29, §1.1].

Now let K be a profinite group. If S is a K-set, then S is a *discrete K-set* if the action map $K \times S \to S$ is continuous, where S is given the discrete topology. Then, a *discrete K-spectrum* Y is a K-spectrum of simplicial sets, such that each simplicial set Y_k is a simplicial discrete K-set (that is, for each $l \geq 0$, $Y_{k,l}$ is a discrete K-set, and all the face and degeneracy maps are K-equivariant). Then, due to work of Jardine and Thomason, as explained in Sections 5 and 7, there is a homotopy fixed point spectrum Y^{hK} defined with respect to the continuous action of K, and, in nice situations, a descent spectral sequence

$$H_c^s(K; \pi_t(Y)) \Rightarrow \pi_{t-s}(Y^{hK}),$$

where the E_2-term is the continuous cohomology of K with coefficients in the discrete K-module $\pi_t(Y)$.

Notice that we use the notation $E_n^{\mathbf{h}G}$ for the construction of Devinatz and Hopkins, and $(-)^{hK}$ for homotopy fixed points with respect to a continuous action, although henceforth, when K is finite and Y is a K-spectrum of topological spaces, we write $Y^{h'K}$ for $\mathrm{holim}_K\, Y$, which is an equivalent definition of the homotopy fixed point spectrum $\mathrm{Map}_K(EK_+, Y)$.

After comparing the spectral sequence for $E_n^{\mathbf{h}G} \wedge X$ with the descent spectral sequence for Y^{hK}, $E_n \wedge X$ appears to be a continuous G_n-spectrum with "descent" spectral sequences for "homotopy fixed point spectra" $E_n^{\mathbf{h}G} \wedge X$. Indeed, we apply [6] to show that $E_n \wedge X$ is a continuous G_n-spectrum; that is, $E_n \wedge X$ is the homotopy limit of a tower of fibrant discrete G_n-spectra. Using this continuous action, we define the homotopy fixed point spectrum $(E_n \wedge X)^{hG}$ and construct its descent spectral sequence.

In more detail, the $K(n)$-local spectrum E_n has an action by the profinite group G_n through maps of commutative S-algebras. The spectrum $E_n^{\mathbf{h}G}$, a $K(n)$-local commutative S-algebra, is referred to as a "homotopy fixed point spectrum" because it has the following desired properties: (a) for any finite spectrum X, there exists spectral sequence (1.2), which has the form of a descent spectral sequence; (b) when G is finite, there is a weak equivalence $E_n^{\mathbf{h}G} \to E_n^{h'G}$, and the descent spectral sequence for $E_n^{h'G}$ is isomorphic to spectral sequence (1.2) (when $X = S^0$) [6, Thm. 3]; and (c) $E_n^{\mathbf{h}G}$ is an $N(G)/G$-spectrum, where $N(G)$ is the normalizer of G in G_n [6, pg. 5]. Thus, their constructions strongly suggest that G_n acts on E_n in a continuous sense.

However, in [6], the action of G_n on E_n is not proven to be continuous, and $E_n^{\mathbf{h}G}$ is not defined with respect to a continuous G-action. Also, when G is profinite, homotopy fixed points should always be the total right derived functor of fixed points, in some sense, and, in [6], it is not shown that the "homotopy fixed point spectrum" $E_n^{\mathbf{h}G}$ can be obtained through such a total right derived functor.

After introducing some notation, we state the main results of this paper. Let BP be the Brown-Peterson spectrum with $BP_* = \mathbb{Z}_{(p)}[v_1, v_2, ...]$, where the degree of v_i is $2(p^i - 1)$. The ideal $(p^{i_0}, v_1^{i_1}, ..., v_{n-1}^{i_{n-1}}) \subset BP_*$ is denoted by I; M_I is the corresponding generalized Moore spectrum $M(p^{i_0}, v_1^{i_1}, ..., v_{n-1}^{i_{n-1}})$, a trivial G_n-spectrum. Given an ideal I, M_I need not exist; however, enough exist for our constructions. The spectrum M_I is a finite type n spectrum with $BP_*(M_I) \cong BP_*/I$. The set $\{i_0, ..., i_{n-1}\}$ of superscripts varies so that there is a family of ideals $\{I\}$. ([4, §4], [19, §4], and [27, Prop. 3.7] provide details for our statements about the spectra M_I.) The map $r\colon BP_* \to E_{n*}$ - defined by $r(v_i) = u_i u^{1-p^i}$, where $u_n = 1$ and $u_i = 0$, when $i > n$ - makes E_{n*} a BP_*-module. Then, by the Landweber exact functor theorem for BP, $\pi_*(E_n \wedge M_I) \cong E_{n*}/I$.

The collection $\{I\}$ contains a descending chain of ideals $\{I_0 \supset I_1 \supset I_2 \supset \cdots\}$, such that there exists a corresponding tower of generalized Moore spectra $\{M_{I_0} \leftarrow M_{I_1} \leftarrow M_{I_2} \leftarrow \cdots\}$. In what follows, the functors $\lim_I$ and holim_I are always taken over the tower of ideals $\{I_i\}$, so that $\lim_I$ and holim_I are really $\lim_{I_i}$ and $\operatorname{holim}_{I_i}$, respectively. Also, in this paper, the homotopy limit of spectra, holim, is constructed levelwise in $\mathcal{S}$, the category of simplicial sets, as defined in [3] and [41, 5.6].

As in [6, (1.4)], let $G_n = U_0 \gneq U_1 \gneq \cdots \gneq U_i \gneq \cdots$ be a descending chain of open normal subgroups, such that $\bigcap_i U_i = \{e\}$ and the canonical map $G_n \to \lim_i G_n/U_i$ is a homeomorphism. We define

$$F_n = \operatorname{colim}_i E_n^{\mathrm{h}U_i}.$$

Then the key to getting our work started is knowing that

$$E_n \wedge M_I \simeq F_n \wedge M_I,$$

and thus, $E_n \wedge M_I$ has the homotopy type of the discrete G_n-spectrum $F_n \wedge M_I$. This result (Corollary 6.5) is not difficult, thanks to the work of Devinatz and Hopkins.

Given a tower $\{Z_I\}$ of discrete G_n-spectra, there is a tower $\{(Z_I)_f\}$, with G_n-equivariant maps $Z_I \to (Z_I)_f$ that are weak equivalences, and $(Z_I)_f$ is a fibrant discrete G_n-spectrum (see Def. 4.1). For the remainder of this section, X is any spectrum with trivial G_n-action. We use $\cong$ to denote an isomorphism in the stable homotopy category.

Theorem 1.3. *As the homotopy limit of a tower of fibrant discrete G_n-spectra, $E_n \cong \operatorname{holim}_I (F_n \wedge M_I)_f$ is a continuous G_n-spectrum. Also, for any spectrum X, $\hat{L}(E_n \wedge X) \cong \operatorname{holim}_I (F_n \wedge M_I \wedge X)_f$ is a continuous G_n-spectrum.*

Using the hypercohomology spectra of Thomason [41], and the theory of presheaves of spectra on a site and globally fibrant models, developed by Jardine (e.g. [21], [22], [23], [24]), we present the theory of homotopy fixed points for discrete G-spectra by considering the site of finite discrete G-sets. Using the fact that G_n has finite virtual cohomological dimension, Thomason's hypercohomology spectrum gives a concrete model for the homotopy fixed points that makes building the descent spectral sequence easy. We apply this theory to define homotopy fixed points for towers of discrete G-spectra.

We point out that much of the theory described above (in Sections 3, 5, 7, and 8, through Remark 8.8) is already known, in some form, especially in the work of Jardine mentioned above, in the excellent article [29], by Mitchell (see also the opening remark of [31, §5]), and in Goerss's paper [11]. However, since the above theory has not been explained in detail before, using the language of homotopy fixed points for discrete G-spectra, we give a presentation of it.

After defining homotopy fixed points for towers of discrete G-spectra, we show that these homotopy fixed points are the total right derived functor of fixed points in the appropriate sense, and we construct the associated descent spectral sequence. This enables us to define the homotopy fixed point spectrum $(\hat{L}(E_n \wedge X))^{hG}$, using the continuous G_n-action, and construct its descent spectral sequence. More specifically, we have the following results.

Definition 1.4. Given a profinite group G, let $\mathcal{O}_G$ be the *orbit category* of G. The objects of $\mathcal{O}_G$ are the continuous left G-spaces G/K, for all K closed in G, and the morphisms are the continuous G-equivariant maps. Note that each object in $\mathcal{O}_G$ is a profinite space.

We use Sp to denote the model category $(\text{spectra})^{\text{stable}}$ of Bousfield-Friedlander spectra.

Theorem 1.5. *There is a functor*

$$P\colon (\mathcal{O}_{G_n})^{\mathrm{op}} \to Sp, \quad P(G_n/G) = E_n^{hG},$$

where G is any closed subgroup of G_n.

We also show that the G-homotopy fixed points of $\hat{L}(E_n \wedge X)$ can be obtained by taking the $K(n)$-localization of the G-homotopy fixed points of the discrete G-spectrum $(F_n \wedge X)$. This result shows that the spectrum F_n is an interesting spectrum that is worth further study.

Theorem 1.6. *For any closed subgroup G and any spectrum X, there is an isomorphism*

$$(\hat{L}(E_n \wedge X))^{hG} \cong \hat{L}((F_n \wedge X)^{hG})$$

in the stable homotopy category. In particular, $E_n^{hG} \cong \hat{L}(F_n^{hG})$.

Theorem 1.7. *Let G be a closed subgroup of G_n and let X be any spectrum. Then there is a conditionally convergent descent spectral sequence*

$$E_2^{s,t} \Rightarrow \pi_{t-s}((\hat{L}(E_n \wedge X))^{hG}). \tag{1.8}$$

If the tower of abelian groups $\{\pi_t(E_n \wedge M_I \wedge X)\}_I$ satisfies the Mittag-Leffler condition, for each $t \in \mathbb{Z}$, then

$$E_2^{s,t} \cong H^s_{\mathrm{cts}}(G; \pi_t(\hat{L}(E_n \wedge X))),$$

the cohomology of continuous cochains. If X is a finite spectrum, then (1.8) has the form

$$H^s_c(G; \pi_t(E_n \wedge X)) \Rightarrow \pi_{t-s}((E_n \wedge X)^{hG}), \tag{1.9}$$

where the E_2-term is the continuous cohomology of (1.2).

Also, Theorem 9.9 shows that, when X is finite, $(E_n \wedge X)^{hG} \cong E_n^{hG} \wedge X$, so that descent spectral sequence (1.9) has the same form as spectral sequence (1.2). It is natural to wonder if these two spectral sequences are isomorphic to each other. Also, the spectra $E_n^{\mathrm{h}G}$ and E_n^{hG} should be the same. We plan to say more about the relationship between $E_n^{\mathrm{h}G}$ and E_n^{hG} and their associated spectral sequences in future work.

It is important to note that the model given here for E_n as a continuous G_n-spectrum is not completely satisfactory, since the continuous action is by morphisms that are just maps of spectra. Because there are models for E_n where the G_n-action is by A_∞- and E_∞-maps of ring spectra, we would like to know that such structured actions are actually continuous.

We outline the contents of this paper. In Section 2, after establishing some notation and terminology, we provide some background material and recall useful facts. In Section 3, we study the model category of discrete G-spectra. In Section 4, we study towers of discrete G-spectra and we give a definition of continuous G-spectrum. In Section 5, we define homotopy fixed points for discrete G-spectra and state some basic facts about this concept. Section 6 shows that E_n is a continuous G_n-spectrum, proving the first half of Theorem 1.3. Section 7 constructs two useful models of the G-homotopy fixed point spectrum, when G has finite virtual cohomological dimension. Section 8 defines homotopy fixed points for towers of discrete G-spectra, builds a descent spectral sequence in this setting, and shows that these homotopy fixed points are a total right derived functor, in the appropriate sense. Section 9 completes the proof of Theorem 1.3, studies $(\hat{L}(E_n \wedge X))^{hG}$, and proves Theorems 1.5 and 1.6. Section 10 considers the descent spectral sequence for $(\hat{L}(E_n \wedge X))^{hG}$ and proves Theorem 1.7.

Acknowledgements. This paper is a development of part of my thesis. I am very grateful to my thesis advisor, Paul Goerss, for many helpful conversations and useful suggestions regarding this paper. Also, I thank Ethan Devinatz for very helpful answers to my questions about his work [6] with Mike Hopkins. I am grateful to Halvard Fausk, Christian Haesemeyer, Rick Jardine, and Charles Rezk for useful conversations.

2. Notation, Terminology, and Preliminaries

We begin by establishing some notation and terminology that will be used throughout the paper. **Ab** is the category of abelian groups. Outside of **Ab**, all groups are assumed to be profinite, unless stated otherwise. For a group G, we write $G \cong \lim_N G/N$, the inverse limit over the open normal subgroups. The notation $H <_c G$ means that H is a closed subgroup of G. We use G to denote arbitrary profinite groups and, specifically, closed subgroups of G_n.

Let $\mathcal{C}$ be a category. A tower $\{C_i\}$ of objects in $\mathcal{C}$ is a diagram in $\mathcal{C}$ of the form $\cdots \to C_i \to C_{i-1} \to \cdots \to C_1 \to C_0$. We always use Bousfield-Friedlander spectra [2], except when another category of spectra is specified. If $\mathcal{C}$ is a model category, then $\mathrm{Ho}(\mathcal{C})$ is its homotopy category. The phrase "stable category" always refers to $\mathrm{Ho}(Sp)$.

In $\mathcal{S}$, the category of simplicial sets, $S^n = \Delta^n/\partial\Delta^n$ is the n-sphere. Given a spectrum X, $X^{(0)} = S^0$, and for $j \geq 1$, $X^{(j)} = X \wedge X \wedge \cdots \wedge X$, with j factors. L_n denotes Bousfield localization with respect to $E(n)_* = \mathbb{Z}_{(p)}[v_1, ..., v_{n-1}][v_n, v_n^{-1}]$.

Definition 2.1. [17, Def. 1.3.1] Let $\mathcal{C}$ and $\mathcal{D}$ be model categories. The functor $F\colon \mathcal{C} \to \mathcal{D}$ is a *left Quillen functor* if F is a left adjoint that preserves cofibrations and trivial cofibrations. The functor $P\colon \mathcal{D} \to \mathcal{C}$ is a *right Quillen functor* if P is a right adjoint that preserves fibrations and trivial fibrations. Also, if F and P are an adjoint pair and left and right Quillen functors, respectively, then (F, P) is a *Quillen pair* for the model categories $(\mathcal{C}, \mathcal{D})$.

Recall [17, Lemma 1.3.10] that a Quillen pair (F, P) yields total left and right derived functors $\mathbf{L}F$ and $\mathbf{R}P$, respectively, which give an adjunction between the homotopy categories $\mathrm{Ho}(\mathcal{C})$ and $\mathrm{Ho}(\mathcal{D})$.

We use $\mathrm{Map}_c(G, A) = \Gamma_G(A)$ to denote the set of continuous maps from G to the topological space A, where A is often a set, equipped with the discrete topology, or a discrete abelian group. Instead of $\Gamma_G(A)$, sometimes we write just $\Gamma(A)$, when the G is understood from context. Let $(\Gamma_G)^k(A)$ denote $(\Gamma_G \Gamma_G \cdots \Gamma_G)(A)$, the application of Γ_G to A, iteratively, $k+1$ times, where $k \geq 0$. Let G^k be the k-fold product of G and let $G^0 = *$. Then, if A is a discrete set or a discrete abelian group, there is a G-equivariant isomorphism $(\Gamma_G)^k(A) \cong \mathrm{Map}_c(G^{k+1}, A)$ of discrete G-sets (modules), where $\mathrm{Map}_c(G^{k+1}, M)$ has G-action defined by

$$(g' \cdot f)(g_1, ..., g_{k+1}) = f(g_1 g', g_2, g_3, ..., g_{k+1}).$$

Also, we often write $\Gamma^k_G(A)$, or $\Gamma^k A$, for $\mathrm{Map}_c(G^k, A)$.

Let A be a discrete abelian group. Then $\mathrm{Map}^\ell_c(G^k_n, A)$ is the discrete G_n-module of continuous maps $G^k_n \to A$ with action defined by

$$(g' \cdot f)(g_1, ..., g_k) = f((g')^{-1} g_1, g_2, g_3, ..., g_k).$$

It is helpful to note that there is a G_n-equivariant isomorphism of discrete G_n-modules

$$p \colon \mathrm{Map}^\ell_c(G^k_n, A) \to \mathrm{Map}_c(G^k_n, A), \quad p(f)(g_1, g_2, ..., g_k) = f(g_1^{-1}, g_2, ..., g_k).$$

$\mathrm{Map}^\ell_c(G^k_n, A)$ is also defined when A is an inverse limit of discrete abelian groups.

By a *topological G-module*, we mean an abelian Hausdorff topological group that is a G-module, with a continuous G-action. Note that if $M = \lim_i M_i$ is the inverse limit of a tower $\{M_i\}$ of discrete G-modules, then M is a topological G-module.

For the remainder of this section, we recall some frequently used facts and discuss background material, to help get our work started.

In [6], Devinatz and Hopkins, using work by Goerss and Hopkins ([12], [13]), and Hopkins and Miller [34], show that the action of G_n on E_n is by maps of commutative S-algebras. Previously, Hopkins and Miller had shown that G_n acts on E_n by maps of A_∞-ring spectra. However, the continuous action presented here is not structured. As already mentioned, the starting point for the continuous action is the spectrum $F_n \wedge M_I$, which is not known to be an A_∞-ring object in the category of discrete G_n-spectra. Thus, we work in the unstructured category Sp of Bousfield-Friedlander spectra of simplicial sets, and the continuous action is simply by maps of spectra.

As mentioned above, [6] is written using E_∞, the category of commutative S-algebras, and $\mathcal{M}_S$, the category of S-modules (see [8]). However, [18, §4.2], [28, §14, §19], and [36, pp. 529-530] show that $\mathcal{M}_S$ and Sp are Quillen equivalent model categories [17, §1.3.3]. Thus, we can import the results of Devinatz and Hopkins from $\mathcal{M}_S$ into Sp. For example, [6, Thm. 1] implies the following result, where $R^+_{G_n}$ is the category whose objects are finite discrete left G_n-sets and G_n itself (a continuous profinite left G_n-space), and whose morphisms are continuous G_n-equivariant maps.

Theorem 2.2 (Devinatz, Hopkins)**.** *There is a presheaf of spectra*

$$F \colon (R^+_{G_n})^{\mathrm{op}} \to Sp,$$

such that (a) for each $S \in R^+_{G_n}$, $F(S)$ *is* $K(n)$*-local; (b)* $F(G_n) = E_n$*; (c) for* U *an open subgroup of* G_n, $E_n^{\mathtt{h}U} := F(G_n/U)$*; and (d)* $F(*) = E_n^{\mathtt{h}G_n} \simeq \hat{L}S^0$.

Now we define a spectrum that is essential to our constructions.

Definition 2.3. Let $F_n = \operatorname{colim}_i E_n^{\mathtt{h}U_i}$, where the direct limit is in Sp. Because $\operatorname{Hom}_{G_n}(G_n/U_i, G_n/U_i) \cong G_n/U_i$, F makes $E_n^{\mathtt{h}U_i}$ a G_n/U_i-spectrum. Thus, F_n is a G_n-spectrum, and the canonical map $\eta\colon F_n \to E_n$ is G_n-equivariant.

Note that F_n is the stalk of the presheaf of spectra $F|_{(G\text{–}\mathbf{Sets}_{df})^{\mathrm{op}}}$, at the unique point of the Grothendieck topos (see §3). The following useful fact is stated in [6, pg. 9] (see also [38, Lemma 14]).

Theorem 2.4. *For* $j \geq 0$, *let* X *be a finite spectrum and regard* $\hat{L}(E_n^{(j+1)} \wedge X)$ *as a* G_n*-spectrum, where* G_n *acts only on the leftmost factor of the smash product. Then there is a* G_n*-equivariant isomorphism*

$$\pi_*(\hat{L}(E_n^{(j+1)} \wedge X)) \cong \operatorname{Map}^{\ell}_c(G_n^j, \pi_*(E_n \wedge X)).$$

We review some frequently used facts about the functor L_n and homotopy limits of spectra. First, L_n is smashing, e.g. $L_nX \simeq X \wedge L_nS^0$, for any spectrum X, and $E(n)$-localization commutes with homotopy direct limits [33, Thms. 7.5.6, 8.2.2]. Note that this implies that F_n is $E(n)$-local.

Definition 2.5. If $\cdots \to X_i \to X_{i-1} \to \cdots \to X_1 \to X_0$ is a tower of spectra such that each X_i is fibrant in Sp, then $\{X_i\}$ is a *tower of fibrant spectra.*

If $\{X_i\}$ is a tower of fibrant spectra, then there is a short exact sequence

$$0 \to \lim{}^1{}_i\, \pi_{m+1}(X_i) \to \pi_m(\operatorname{holim}_i X_i) \to \lim_i \pi_m(X_i) \to 0.$$

Also, if each map in the tower is a fibration, the map $\lim_i X_i \to \operatorname{holim}_i X_i$ is a weak equivalence. If J is a small category and the functor $P\colon J \to Sp$ is a diagram of spectra, such that P_j is fibrant for each $j \in J$, then $\operatorname{holim}_j P_j$ is a fibrant spectrum.

Definition 2.6. There is a functor $(-)_{\mathtt{f}}\colon Sp \to Sp$, such that, given Y in Sp, $Y_{\mathtt{f}}$ is a fibrant spectrum, and there is a natural transformation $\mathrm{id}_{Sp} \to (-)_{\mathtt{f}}$, such that, for any Y, the map $Y \to Y_{\mathtt{f}}$ is a trivial cofibration. For example, if Y is a G-spectrum, then $Y_{\mathtt{f}}$ is also a G-spectrum, and the map $Y \to Y_{\mathtt{f}}$ is G-equivariant.

The following statement says that smashing with a finite spectrum commutes with homotopy limits.

Lemma 2.7 ([42, pg. 96])**.** *Let* J *be a small category,* $\{Z_j\}$ *a* J*-shaped diagram of fibrant spectra, and let* Y *be a finite spectrum. Then the composition*

$$(\operatorname{holim}_j Z_j) \wedge Y \to \operatorname{holim}_j (Z_j \wedge Y) \to \operatorname{holim}_j (Z_j \wedge Y)_{\mathtt{f}}$$

is a weak equivalence.

We recall the result that is used to build towers of discrete G-spectra.

Theorem 2.8 ([16, §2], [5, Remark 3.6])**.** *If* X *is an* $E(n)$*-local spectrum, then, in the stable category, there is an isomorphism*

$$\hat{L}X \cong \operatorname{holim}_I (X \wedge M_I)_{\mathtt{f}}.$$

Lemma 2.9 ([19, Lemma 7.2])**.** *If* X *is any spectrum, and* Y *is a finite spectrum of type* n, *then* $\hat{L}(X \wedge Y) \simeq \hat{L}(X) \wedge Y \simeq L_n(X) \wedge Y$.

We recall some useful facts about compact p-adic analytic groups. Since S_n is compact p-adic analytic, and G_n is an extension of S_n by the Galois group, G_n is a compact p-adic analytic group [37, Cor. of Thm. 2]. Any closed subgroup of a compact p-adic analytic group is also compact p-adic analytic [7, Thm. 9.6]. Also, since the subgroup in S_n of strict automorphisms is finitely generated and pro-p, [35, pp. 76, 124] implies that all subgroups in G_n of finite index are open.

Let the profinite group G be a compact p-adic analytic group. Then G contains an open subgroup H, such that H is a pro-p group with finite cohomological p-dimension; that is, $\mathrm{cd}_p(H) = m$, for some non-negative integer m (see [25, 2.4.9] or the exposition in [39]). Since H is pro-p, $\mathrm{cd}_q(H) = 0$, whenever q is a prime different from p [45, Prop. 11.1.4]. Also, if M is a discrete H-module, then, for $s \geq 1$, $H^s_c(H; M)$ is a torsion abelian group [35, Cor. 6.7.4]. These facts imply that, for any discrete H-module M, $H^s_c(H; M) = 0$, whenever $s > m + 1$. We express this conclusion by saying that G has finite virtual cohomological dimension and we write $\mathrm{vcd}(G) \leq m$. Also, if K is a closed subgroup of G, $H \cap K$ is an open pro-p subgroup of K with $\mathrm{cd}_p(H \cap K) \leq m$, so that $\mathrm{vcd}(K) \leq m$, and thus, m is a uniform bound independent of K.

Now we state various results related to towers of abelian groups and continuous cohomology. The lemma below follows from the fact that a tower of abelian groups satisfies the Mittag-Leffler condition if and only if the tower is pro-isomorphic to a tower of epimorphisms [20, (1.14)].

Lemma 2.10. *Let $F\colon \mathbf{Ab} \to \mathbf{Ab}$ be an exact additive functor. If $\{A_i\}_{i\geq 0}$ is a tower of abelian groups that satisfies the Mittag-Leffler condition, then so does the tower $\{F(A_i)\}$.*

Remark 2.11. Let G be a profinite group. Then the functor

$$\mathrm{Map}_c(G, -)\colon \mathbf{Ab} \to \mathbf{Ab}, \quad A \mapsto \mathrm{Map}_c(G, A),$$

is defined by giving A the discrete topology. The isomorphism $\mathrm{Map}_c(G, A) \cong \operatorname{colim}_N \prod_{G/N} A$ shows that $\mathrm{Map}_c(G, -)$ is an exact additive functor. Later, we will use Lemma 2.10 with this functor.

The next lemma is a consequence of the fact that limits in $\mathbf{Ab}$ and in topological spaces are created in $\mathbf{Sets}$.

Lemma 2.12. *Let $M = \lim_\alpha M_\alpha$ be an inverse limit of discrete abelian groups, so that M is an abelian topological group. Let H be any profinite group. Then $\mathrm{Map}_c(H, M) \to \lim_\alpha \mathrm{Map}_c(H, M_\alpha)$ is an isomorphism of abelian groups.*

Lemma 2.13. *If X is a finite spectrum, $G <_c G_n$, and t any integer, then the abelian group $\pi_t(E_n \wedge M_I \wedge X)$ is finite.*

Proof. The starting point is the fact that $\pi_t(E_n \wedge M_I) \cong \pi_t(E_n)/I$ is finite. In the stable homotopy category of CW-spectra, since X is a finite spectrum, there exists some m such that $X_l = \Sigma^{l-m} X_m$ whenever $l \geq m$, and X_m is a finite complex. Since $\pi_t(E_n \wedge M_I \wedge X) \cong \pi_{t-m}(E_n \wedge M_I \wedge X_m)$ and X_m can be built out of a finite number of cofiber sequences, the result follows. □

Corollary 2.14 ([15, pg. 116]). *If X is a finite spectrum, then $\pi_t(E_n \wedge X) \cong \lim_I \pi_t(E_n \wedge M_I \wedge X)$.*

We recall the definition of a second version of continuous cohomology. If M is a topological G-module, then $H^s_{\mathrm{cts}}(G; M)$ is the sth cohomology group of the cochain complex $M \to \mathrm{Map}_c(G, M) \to \mathrm{Map}_c(G^2, M) \to \cdots$ of continuous cochains for a profinite group G with coefficients in M (see [40, §2]). If M is a discrete G-module, this is the usual continuous cohomology $H^s_c(G; M)$. There is the following useful relationship between these cohomology theories.

Theorem 2.15 ([20, (2.1), Thm. 2.2])**.** *Let $\{M_n\}_{n\geq 0}$ be a tower of discrete G-modules satisfying the Mittag-Leffler condition and let $M = \lim_n M_n$ as a topological G-module. Then, for each $s \geq 0$, there is a short exact sequence*

$$0 \to {\lim}^1{}_n\, H^{s-1}_c(G; M_n) \to H^s_{\mathrm{cts}}(G; M) \to \lim{}_n\, H^s_c(G; M_n) \to 0,$$

where $H^{-1}_c(G; -) = 0$.

The next result is from [6, Rk. 1.3] and is due to the fact that, when X is a finite spectrum, $\pi_t(E_n \wedge X) \cong \lim_k \pi_t(E_n \wedge X)/I^k_n \pi_t(E_n \wedge X)$ is a profinite continuous $\mathbb{Z}_p[\![G]\!]$-module.

Lemma 2.16. *If G is closed in G_n and X is a finite spectrum, then, for $s \geq 0$,*

$$\begin{aligned} H^s_{\mathrm{cts}}(G; \pi_t(E_n \wedge X)) &\cong \lim{}_k\, H^s_c(G; \pi_t(E_n \wedge X)/I^k_n \pi_t(E_n \wedge X)) \\ &= H^s_c(G; \pi_t(E_n \wedge X)). \end{aligned}$$

We make a few remarks about the functorial smash product in *Sp*, defined in [23, Chps. 1, 2].

Definition 2.17. Given spectra X and Y, their smash product $X \wedge Y$ is given by $(X \wedge Y)_{2k} = X_k \wedge Y_k$ and $(X \wedge Y)_{2k+1} = X_k \wedge Y_{k+1}$, where, for example, $X_k \in \mathcal{S}_*$, the category of pointed simplicial sets.

Since $K \wedge (-) \colon \mathcal{S}_* \to \mathcal{S}_*$ is a left adjoint, for any K in $\mathcal{S}_*$, smashing with any spectrum in either variable commutes with colimits in *Sp*.

3. The model category of discrete G-spectra

A pointed simplicial discrete G-set is a pointed simplicial set that is a simplicial discrete G-set, such that the G-action fixes the basepoint.

Definition 3.1. A *discrete G-spectrum* X is a spectrum of pointed simplicial sets X_k, for $k \geq 0$, such that each simplicial set X_k is a pointed simplicial discrete G-set, and each bonding map $S^1 \wedge X_k \to X_{k+1}$ is G-equivariant (S^1 has trivial G-action). Let Sp_G denote the category of discrete G-spectra, where the morphisms are G-equivariant maps of spectra.

As with discrete G-sets, if $X \in Sp_G$, there is a G-equivariant isomorphism $X \cong \mathrm{colim}_N\, X^N$. Also, a discrete G-spectrum X is a continuous G-spectrum since, for all $k, l \geq 0$, the set $X_{k,l}$ is a continuous G-space with the discrete topology, and all the face and degeneracy maps are (trivially) continuous.

Definition 3.2. As in [23, §6.2], let $G-\mathbf{Sets}_{df}$ be the canonical site of finite discrete G-sets. The pretopology of $G-\mathbf{Sets}_{df}$ is given by covering families of the form $\{f_\alpha \colon S_\alpha \to S\}$, a finite set of G-equivariant functions in $G-\mathbf{Sets}_{df}$ for a fixed $S \in G-\mathbf{Sets}_{df}$, such that $\coprod_\alpha S_\alpha \to S$ is a surjection.

We use **Shv** to denote the category of sheaves of sets on the site $G-\mathbf{Sets}_{df}$. Also, T_G signifies the category of discrete G-sets, and, as in [11], S_G is the category of simplicial objects in T_G. The Grothendieck topos **Shv** has a unique point $u\colon \mathbf{Sets} \to \mathbf{Shv}$. The left adjoint of the topos morphism u is

$$u^*\colon \mathbf{Shv} \to \mathbf{Sets}, \quad \mathcal{F} \mapsto \operatorname{colim}_N \mathcal{F}(G/N),$$

with right adjoint

$$u_*\colon \mathbf{Sets} \to \mathbf{Shv}, \quad X \mapsto \operatorname{Hom}_G(-, \operatorname{Map}_c(G, X))$$

[23, Rk. 6.25]. The G-action on the discrete G-set $\operatorname{Map}_c(G, X)$ is defined by $(g \cdot f)(g') = f(g'g)$, for g, g' in G, and f a continuous map $G \to X$, where X is given the discrete topology.

The functor $\operatorname{Map}_c(G, -)\colon \mathbf{Sets} \to T_G$ prolongs to the functor

$$\operatorname{Map}_c(G, -)\colon Sp \to Sp_G.$$

Thus, if X is a spectrum, then $\operatorname{Map}_c(G, X) \cong \operatorname{colim}_N \prod_{G/N} X$ is the discrete G-spectrum with $(\operatorname{Map}_c(G, X))_k = \operatorname{Map}_c(G, X_k)$, where $\operatorname{Map}_c(G, X_k)$ is a pointed simplicial set, with l-simplices $\operatorname{Map}_c(G, X_{k,l})$ and basepoint $G \to *$, where $X_{k,l}$ is regarded as a discrete set. The G-action on $\operatorname{Map}_c(G, X)$ is defined by the G-action on the sets $\operatorname{Map}_c(G, X_{k,l})$.

It is not hard to see that $\operatorname{Map}_c(G, -)$ is right adjoint to the forgetful functor $U\colon Sp_G \to Sp$. Note that if X is a discrete G-spectrum, then there is a contravariant functor (presheaf) $\operatorname{Hom}_G(-, X)\colon (G\text{–}\mathbf{Sets}_{df})^{\mathrm{op}} \to Sp$, where, for any $S \in G\text{–}\mathbf{Sets}_{df}$, $\operatorname{Hom}_G(S, X)$ is the spectrum with $(\operatorname{Hom}_G(S, X))_{k,l} = \operatorname{Hom}_G(S, X_{k,l})$, a pointed set with basepoint $S \to *$.

Let **ShvSpt** be the category of sheaves of spectra on the site $G\text{–}\mathbf{Sets}_{df}$. A sheaf of spectra $\mathcal{F}$ is a presheaf $\mathcal{F}\colon (G\text{–}\mathbf{Sets}_{df})^{\mathrm{op}} \to Sp$, such that, for any $S \in G\text{–}\mathbf{Sets}_{df}$ and any covering family $\{f_\alpha\colon S_\alpha \to S\}$, the usual diagram (of spectra) is an equalizer. Equivalently, a sheaf of spectra $\mathcal{F}$ consists of pointed simplicial sheaves $\mathcal{F}^n$, together with pointed maps of simplicial presheaves $\sigma\colon S^1 \wedge \mathcal{F}^n \to \mathcal{F}^{n+1}$, for $n \geq 0$, where S^1 is the constant simplicial presheaf. A morphism between sheaves of spectra is a natural transformation between the underlying presheaves.

The category **PreSpt** of presheaves of spectra on the site $G-\mathbf{Sets}_{df}$ has the following "stable" model category structure ([22], [23, §2.3]). A map $h\colon \mathcal{F} \to \mathcal{G}$ of presheaves of spectra is a weak equivalence if and only if the associated map of stalks $\operatorname{colim}_N \mathcal{F}(G/N) \to \operatorname{colim}_N \mathcal{G}(G/N)$ is a weak equivalence of spectra. Recall that a map k of simplicial presheaves is a cofibration if, for each $S \in G-\mathbf{Sets}_{df}$, $k(S)$ is a monomorphism of simplicial sets. Then h is a cofibration of presheaves of spectra if the following two conditions hold:

(1) the map $h^0\colon \mathcal{F}^0 \to \mathcal{G}^0$ is a cofibration of simplicial presheaves; and
(2) for each $n \geq 0$, the canonical map $(S^1 \wedge \mathcal{G}^n) \cup_{S^1 \wedge \mathcal{F}^n} \mathcal{F}^{n+1} \to \mathcal{G}^{n+1}$ is a cofibration of simplicial presheaves.

Fibrations are those maps with the right lifting property with respect to trivial cofibrations.

Definition 3.3. In the stable model category structure, fibrant presheaves are often referred to as *globally fibrant*, and if $\mathcal{F} \to \mathcal{G}$ is a weak equivalence of presheaves, with $\mathcal{G}$ globally fibrant, then $\mathcal{G}$ is a *globally fibrant model* for $\mathcal{F}$. We often use $\mathbf{G}\mathcal{F}$ to denote such a globally fibrant model.

We recall the following fact, which is especially useful when $S = *$.

Lemma 3.4. *Let $S \in G-\mathbf{Sets}_{df}$. The S-sections functor $\mathbf{PreSpt} \to Sp$, defined by $\mathcal{F} \mapsto \mathcal{F}(S)$, preserves fibrations, trivial fibrations, and weak equivalences between fibrant objects.*

Proof. The S-sections functor has a left adjoint, obtained by left Kan extension, that preserves cofibrations and weak equivalences. See [23, pg. 60] and [29, Cor. 3.16] for the details. □

We use $\mathcal{L}^2$ to denote the sheafification functor for presheaves of sets, simplicial presheaves, and presheaves of spectra. Let $i\colon \mathbf{ShvSpt} \to \mathbf{PreSpt}$ be the inclusion functor, which is right adjoint to $\mathcal{L}^2$. In [10, Rk. 3.11], $\mathbf{ShvSpt}$ is given the following model category structure. A map $h\colon \mathcal{F} \to \mathcal{G}$ of sheaves of spectra is a weak equivalence (fibration) if and only if $i(f)$ is a weak equivalence (fibration) of presheaves. Also, h is a cofibration of sheaves of spectra if the following two conditions hold:

1. the map $h^0\colon \mathcal{F}^0 \to \mathcal{G}^0$ is a cofibration of simplicial presheaves; and
2. for each $n \geq 0$, the canonical map $\mathcal{L}^2((S^1 \wedge \mathcal{G}^n) \cup_{S^1 \wedge \mathcal{F}^n} \mathcal{F}^{n+1}) \to \mathcal{G}^{n+1}$ is a cofibration of simplicial presheaves.

Since i preserves weak equivalences and fibrations between sheaves of spectra, $\mathcal{L}^2$ preserves cofibrations, trivial cofibrations, and weak equivalences between cofibrant objects, and $(\mathcal{L}^2, i)$ is a Quillen pair for $(\mathbf{PreSpt}, \mathbf{ShvSpt})$. By [23, Cor. 6.22], if $\mathcal{F}$ is a presheaf of sets, a simplicial presheaf, or a presheaf of spectra, then

$$\mathcal{L}^2\mathcal{F} \cong \mathrm{Hom}_G(-, \operatorname{colim}_N \mathcal{F}(G/N)). \tag{3.5}$$

This implies that, for any presheaf of spectra $\mathcal{F}$, $\mathcal{F} \to \mathcal{L}^2\mathcal{F}$ is a weak equivalence, and thus, $\mathrm{Ho}(\mathbf{PreSpt}) \cong \mathrm{Ho}(\mathbf{ShvSpt})$ is a Quillen equivalence.

There is an equivalence of categories $\mathbf{ShvSpt} \cong Sp_G$, via the functors

$$L\colon \mathbf{ShvSpt} \to Sp_G, \quad L(\mathcal{F}) = \operatorname{colim}_N \mathcal{F}(G/N), \quad \text{and}$$

$$R\colon Sp_G \to \mathbf{ShvSpt}, \quad R(X) = \mathrm{Hom}_G(-, X).$$

It is not hard to verify this equivalence, since it is an extension to spectra of the fact that $\mathbf{Shv}$ and T_G are equivalent categories (see [23, Prop. 6.20], [26, III-9, Thm. 1]).

Exploiting this equivalence, we make Sp_G a model category in the following way. Define a map f of discrete G-spectra to be a weak equivalence (fibration) if and only if $\mathrm{Hom}_G(-, f)$ is a weak equivalence (fibration) of sheaves of spectra. Also, define f to be a cofibration if and only if f has the left lifting property with respect to all trivial fibrations. Thus, f is a cofibration if and only if $\mathrm{Hom}_G(-, f)$ is a cofibration in $\mathbf{ShvSpt}$. Using this, it is easy to show that Sp_G is a model category, and there is a Quillen equivalence $\mathrm{Ho}(\mathbf{ShvSpt}) \cong \mathrm{Ho}(Sp_G)$.

Formula (3.5) allows us to define the model category structure of Sp_G without reference to sheaves of spectra, in the theorem below, extending the model category structure on the category S_G, given in [11, Thm. 1.12], to Sp_G.

Theorem 3.6. *Let $f\colon X \to Y$ be a map in Sp_G. Then f is a weak equivalence (cofibration) in Sp_G if and only if f is a weak equivalence (cofibration) in Sp.*

Proof. For weak equivalences, the statement is clearly true. Assume that f is a cofibration in Sp_G. Since $\mathrm{Hom}_G(-, X_0) \to \mathrm{Hom}_G(-, Y_0)$ is a cofibration of simplicial presheaves, evaluation at G/N implies that $X_0^N \to Y_0^N$ is a cofibration of simplicial sets. Thus, $X_0 \cong \mathrm{colim}_N\, X_0^N \to \mathrm{colim}_N\, Y_0^N \cong Y_0$ is a cofibration of simplicial sets.

By (3.5) and the fact that colimits commute with pushouts,

$$\mathrm{Hom}_G(-, (S^1 \wedge Y_n) \cup_{S^1 \wedge X_n} X_{n+1}) \to \mathrm{Hom}_G(-, Y_{n+1})$$

is a cofibration of simplicial presheaves, and hence, the map of simplicial sets $\mathrm{colim}_N\, ((S^1 \wedge Y_n) \cup_{S^1 \wedge X_n} X_{n+1})^N \to Y_{n+1}$ is a cofibration.

Let W be a simplicial pointed discrete G-set. Then $S^1 \wedge W \cong \mathrm{colim}_N\, (S^1 \wedge W^N)$, so that $S^1 \wedge W$ is also a simplicial pointed discrete G-set. Since the forgetful functor $U \colon T_G \to \mathbf{Sets}$ is a left adjoint, pushouts in T_G are formed in $\mathbf{Sets}$, and thus, there is an isomorphism

$$\mathrm{colim}_N\, ((S^1 \wedge Y_n) \cup_{S^1 \wedge X_n} X_{n+1})^N \cong (S^1 \wedge Y_n) \cup_{S^1 \wedge X_n} X_{n+1}$$

of simplicial discrete G-sets. Hence, $(S^1 \wedge Y_n) \cup_{S^1 \wedge X_n} X_{n+1} \to Y_{n+1}$ is a cofibration in $\mathcal{S}$, and f is a cofibration in Sp.

The converse follows from the fact that if $W \to Z$ is an injection of simplicial discrete G-sets, then $\mathrm{Hom}_G(-, W) \to \mathrm{Hom}_G(-, Z)$ is a cofibration of simplicial presheaves. □

Corollary 3.7. *The functors* $(U, \mathrm{Map}_c(G,-))$ *are a Quillen pair for* (Sp_G, Sp)*.*

Let $\mathrm{t} \colon Sp \to Sp_G$ give a spectrum trivial G-action, so that $\mathrm{t}(X) = X$. The right adjoint of t is the fixed points functor $(-)^G$. Clearly, t preserves all weak equivalences and cofibrations, giving the next result.

Corollary 3.8. *The functors* $(\mathrm{t}, (-)^G)$ *are a Quillen pair for* (Sp, Sp_G)*.*

We conclude this section with a few more useful facts about discrete G-spectra.

Lemma 3.9. *If $f \colon X \to Y$ is a fibration in Sp_G, then it is a fibration in Sp. In particular, if X is fibrant as a discrete G-spectrum, then X is fibrant as a spectrum.*

Proof. Since $\mathrm{Hom}_G(-, f)$ is a fibration of presheaves of spectra, $\mathrm{Hom}_G(G/N, f)$ is a fibration of spectra for each open normal subgroup N. This implies that $\mathrm{colim}_N\, \mathrm{Hom}_G(G/N, f)$ is a fibration of spectra. Then the lemma follows from factoring f as $X \cong \mathrm{colim}_N\, X^N \to \mathrm{colim}_N\, Y^N \cong Y$. □

The next lemma and its corollary show that the homotopy groups of a discrete G-spectrum are discrete G-modules, as expected.

Lemma 3.10. *If X is a pointed Kan complex and a simplicial discrete G-set, then $\pi_n(X)$ is a discrete G-module, for all $n \geq 2$.*

Proof. Let $f \colon S^n \to X$ be a pointed map. Since S^n has only two non-degenerate simplices, the basepoint $*$ and the fundamental class $\iota_n \in (S^n)_n$, the map f is determined by $f(\iota_n)$. The action of G on $\pi_n(X)$ is defined as follows: $g \cdot [f]$ is the homotopy class of the pointed map $h \colon S^n \to X$ defined by $h(\iota_n) = g \cdot f(\iota_n)$. To show that the action is continuous, it suffices to show that, for any $[f] \in \pi_n(X)$, the stabilizer $G_{[f]}$ of $[f]$ is an open subgroup of G, and this is elementary. □

Corollary 3.11. *If X is a discrete G-spectrum, then $\pi_n(X)$ is a discrete G-module for any integer n.*

Proof. Let $X \to X_f$ be a trivial cofibration with X_f fibrant, all in Sp_G. Then

$$\pi_n(X) \cong \pi_n(X_f) \cong \operatorname{colim}_{m \geq \max(2-n,0)} \pi_{m+n}((X_f)_m)$$

is a discrete G-module, since each $\pi_{m+n}((X_f)_m)$ is one. □

The following observation says that certain elementary constructions with discrete G-spectra yield discrete G-spectra.

Lemma 3.12. *Given a profinite group $G \cong \lim_N G/N$, let $\{X_N\}_N$ be a directed system of spectra, such that each X_N is a G/N-spectrum and the maps are G-equivariant. Then $\operatorname{colim}_N X_N$ is a discrete G-spectrum. If Y is a trivial G-spectrum, then $(\operatorname{colim}_N X_N) \wedge Y \cong \operatorname{colim}_N (X_N \wedge Y)$ is a G-equivariant isomorphism of discrete G-spectra. Thus, if X is a discrete G-spectrum, then $X \wedge Y$ is a discrete G-spectrum.*

The corollary below is very useful later.

Corollary 3.13. *The spectra F_n, $F_n \wedge M_I$, and $F_n \wedge M_I \wedge X$, for any spectrum X, are discrete G_n-spectra.*

4. Towers of discrete G-spectra and continuous G-spectra

Let $\mathbf{tow}(Sp_G)$ be the category where a typical object $\{X_i\}$ is a tower

$$\cdots \to X_i \to X_{i-1} \to \cdots \to X_1 \to X_0$$

in Sp_G. The morphisms are natural transformations $\{X_i\} \to \{Y_i\}$, such that each $X_i \to Y_i$ is G-equivariant. Since Sp_G is a simplicial model category, [14, VI, Prop. 1.3] shows that $\mathbf{tow}(Sp_G)$ is also a simplicial model category, where $\{f_i\}$ is a weak equivalence (cofibration) if and only if each f_i is a weak equivalence (cofibration) in Sp_G. By [14, VI, Rk. 1.5], if $\{X_i\}$ is fibrant in $\mathbf{tow}(Sp_G)$, then each map $X_i \to X_{i-1}$ in the tower is a fibration and each X_i is fibrant, all in Sp_G.

Definition 4.1. Let $\{X_i\}$ be in $\mathbf{tow}(Sp_G)$. Then $\{X_i'\}$ denotes the target of a trivial cofibration $\{X_i\} \to \{X_i'\}$, with $\{X_i'\}$ fibrant, in $\mathbf{tow}(Sp_G)$.

Remark 4.2. Let $\{X_\alpha\}$ be a diagram in Sp_G. Since there is an isomorphism $\lim_\alpha^{Sp_G} X_\alpha \cong \operatorname{colim}_N (\lim_\alpha^{Sp} X_\alpha)^N$, limits in Sp_G are not formed in Sp. In everything that follows, lim and holim are always in Sp.

The functor $\lim_i (-)^G \colon \mathbf{tow}(Sp_G) \to Sp$, given by $\{X_i\} \mapsto \lim_i X_i^G$, is right adjoint to the functor $\underline{\mathrm{t}} \colon Sp \to \mathbf{tow}(Sp_G)$ that sends a spectrum X to the constant diagram $\{X\}$, where X has trivial G-action. Since $\underline{\mathrm{t}}$ preserves all weak equivalences and cofibrations, we have the following fact.

Lemma 4.3. *The functors $(\underline{\mathrm{t}}, \lim_i (-)^G)$ are a Quillen pair for the categories $(Sp, \mathbf{tow}(Sp_G))$.*

This Lemma implies the existence of the total right derived functor

$$\mathbf{R}(\lim_i (-)^G) \colon \mathrm{Ho}(\mathbf{tow}(Sp_G)) \to \mathrm{Ho}(Sp), \quad \{X_i\} \mapsto \lim_i (X_i')^G.$$

Lemma 4.4. *If $\{X_i\}$ in $\mathbf{tow}(Sp_G)$ is a tower of fibrant spectra, then there are weak equivalences* $\operatorname{holim}_i X_i \xrightarrow{p} \operatorname{holim}_i X_i' \xleftarrow{q} \lim_i X_i'$.

Proof. Since each $X_i \to X_i'$ is a weak equivalence between fibrant spectra, p is a weak equivalence. Since $X_i' \to X_{i-1}'$ is a fibration in Sp, for $i \geq 1$, q is a weak equivalence. □

We use this lemma to define the continuous G-spectra that we will study.

Definition 4.5. If $\{X_i\}$ is in $\mathbf{tow}(Sp_G)$, then the inverse limit $\lim_i X_i$ is a *continuous G-spectrum*. Also, if $\{X_i\} \in \mathbf{tow}(Sp_G)$ is a tower of fibrant spectra, we call $\operatorname{holim}_i X_i$ a *continuous G-spectrum*, due to the zigzag of Lemma 4.4 that relates $\operatorname{holim}_i X_i$ to the continuous G-spectrum $\lim_i X_i'$. Notice that if $X \in Sp_G$, then, using the constant tower on X, $X \cong \lim_i X$ is a continuous G-spectrum.

Remark 4.6. Sometimes we will use the term "continuous G-spectrum" more loosely. Let X be a continuous G-spectrum, as in Definition 4.5. If Y is a G-spectrum that is isomorphic to X, in the stable category, with compatible G-actions, then we call Y a continuous G-spectrum.

We make a few comments about Definition 4.5. Though the definition is not as general as it could be, it is sufficient for our applications. The inverse limit is central to the definition since the inverse limit of a tower of discrete G-sets is a topological G-space.

Given any tower $X_* = \{X_i\}$ in Sp_G,

$$\operatorname{holim}_i X_i = \operatorname{Tot}(\textstyle\prod^* X_*) \cong \lim_n \mathrm{T}(n),$$

where $\mathrm{T}(n) = \operatorname{Tot}_n(\prod^* X_*)$. (See §5 for the definition of $\prod^* X_*$, and [3] is a reference for any undefined notation in this paragraph.) Then it is natural to ask if $T(n)$ is in Sp_G, so that $\operatorname{holim}_i X_i$ is canonically a continuous G-spectrum. For this to be true, it must be that, for any $m \geq 0$, the simplicial set $\mathrm{T}(n)_m = \operatorname{Map}_{c\mathcal{S}}(\mathrm{sk}_n\Delta[-], \prod^*(X_*)_m)$ belongs to $\mathcal{S}_G$. If, for all $s \geq 0$, $\prod^s(X_*)_m \in \mathcal{S}_G$, then $\mathrm{T}(n)_m \in \mathcal{S}_G$, by [11, pg. 212]. However, the infinite product $\prod^s(X_*)_m$ need not be in $\mathcal{S}_G$, and thus, in general, $\mathrm{T}(n) \notin Sp_G$. Therefore, $\operatorname{holim}_i X_i$ is not always identifiable with a continuous G-spectrum, in the above way.

5. Homotopy fixed points for discrete G-spectra

In this section, we define the homotopy fixed point spectrum for $X \in Sp_G$. We begin by recalling the homotopy spectral sequence, since we use it often.

If J is a small category, $P\colon J \to Sp$ a diagram of fibrant spectra, and Z any spectrum, then there is a conditionally convergent spectral sequence

$$E_2^{s,t} = \lim{}^s{}_J\, [Z, P]_t \Rightarrow [Z, \operatorname{holim}_J P]_{t-s}, \tag{5.1}$$

where $\lim^s$ is the sth right derived functor of $\lim_J\colon \mathbf{Ab}^J \to \mathbf{Ab}$ [41, Prop. 5.13, Lem. 5.31].

Associated to P is the cosimplicial spectrum $\prod^* P$, with $\prod^n P = \prod_{(B\Delta)_n} P(j_n)$, where the n-simplices of the nerve $B\Delta$ consist of all strings $[j_0] \to \cdots \to [j_n]$ of n morphisms in Δ. For any $k \geq 0$, $(\prod^* P)_k$ is a fibrant cosimplicial pointed simplicial set, and $\operatorname{holim}_J P = \operatorname{Tot}(\prod^* P)$.

If P is a cosimplicial diagram of fibrant spectra (a *cosimplicial fibrant spectrum*), then $E_2^{s,t} = \pi^s[Z, P]_t$, the sth cohomotopy group of the cosimplicial abelian group $[Z, P]_t$.

Definition 5.2. Given a discrete G-spectrum X, X^{hG} denotes the *homotopy fixed point spectrum* of X with respect to the continuous action of G. We define $X^{hG} = (X_{f,G})^G$, where $X \rightarrow X_{f,G}$ is a trivial cofibration and $X_{f,G}$ is fibrant, in Sp_G. We write X_f instead of $X_{f,G}$ when doing so causes no confusion.

Note that $X^{hG} = \mathrm{Hom}_G(*, X_{f,G})$, the global sections of the presheaf of spectra $\mathrm{Hom}_G(-, X_{f,G})$. This definition has been developed in other categories: see [11] for simplicial discrete G-sets, [21] for simplicial presheaves, and [23, Ch. 6] for presheaves of spectra.

As expected, the definition of homotopy fixed points for a profinite group generalizes the definition for a finite group. Let G be a finite group, X a G-spectrum, and let $X \rightarrow X_{\mathtt{f}}$ be a weak equivalence that is G-equivariant, with $X_{\mathtt{f}}$ a fibrant spectrum. Then, since G is finite, the homotopy fixed point spectrum $X^{h'G} = \mathrm{Map}_G(EG_+, X_{\mathtt{f}})$ can be defined to be $\mathrm{holim}_G\, X_{\mathtt{f}}$ (as in the Introduction). Note that there is a descent spectral sequence

$$E_2^{s,t} = \lim{}^s{}_G\, \pi_t(X) \cong H^s(G; \pi_t(X)) \Rightarrow \pi_{t-s}(X^{h'G}).$$

Since G is profinite, X is a discrete G-spectrum, and $X_{f,G}$ is a fibrant spectrum, so that $X^{h'G} = \mathrm{holim}_G\, X_f$. Then, by [23, Prop. 6.39], the canonical map $X^{hG} = \lim_G X_f \rightarrow \mathrm{holim}_G\, X_f = X^{h'G}$ is a weak equivalence, as desired.

We point out several properties of homotopy fixed points that follow from Corollary 3.8.

Lemma 5.3. *The homotopy fixed points functor* $(-)^{hG} \colon \mathrm{Ho}(Sp_G) \rightarrow \mathrm{Ho}(Sp)$ *is the total right derived functor of the fixed points functor* $(-)^G \colon Sp_G \rightarrow Sp$. *In particular, if* $X \rightarrow Y$ *is a weak equivalence of discrete* G*-spectra, then* $X^{hG} \rightarrow Y^{hG}$ *is a weak equivalence.*

Given two globally fibrant models $\mathbf{G}\mathcal{F}$ and $\mathbf{G}'\mathcal{F}$ for a presheaf of spectra $\mathcal{F}$, there need not be a weak equivalence $\mathbf{G}\mathcal{F} \rightarrow \mathbf{G}'\mathcal{F}$, since not every presheaf of spectra is cofibrant. Also, a cofibration of sheaves of spectra can fail to be a cofibration of presheaves. Given these limitations, there is the following relationship between homotopy fixed points and the global sections of globally fibrant models, due to the left lifting property of trivial cofibrations with respect to fibrations.

Lemma 5.4. *Let* $X \in Sp_G$. *Then* $X^{hG} \xleftarrow{\simeq} \mathcal{F}(*) \xrightarrow{\simeq} \mathbf{G}\mathrm{Hom}_G(-, X)(*)$, *where* $\mathrm{Hom}_G(-, X) \rightarrow \mathcal{F}$ *is a trivial cofibration of presheaves, with* $\mathcal{F}$ *globally fibrant. Thus,* X^{hG} *and* $\mathbf{G}\mathrm{Hom}_G(-, X)(*)$ *have the same stable homotopy type.*

6. E_n IS A CONTINUOUS G_n-SPECTRUM

We show that E_n is a continuous G_n-spectrum by successively eliminating simpler ways of constructing a continuous action, and by applying the theory of the previous section.

First of all, since the profinite ring $\pi_0(E_n)$ is not a discrete G_n-module, Corollary 3.11 implies the following observation.

Lemma 6.1. *E_n is not a discrete G_n-spectrum.*

However, note that, for $k \in \mathbb{Z}$, $\pi_{2k}(E_n \wedge M_I)$ is a finite discrete G_n-module so that the action factors through a finite quotient G_n/U_I, where U_I is some open normal subgroup (see [35, Lem. 1.1.16]). Thus, $\pi_{2k}(E_n \wedge M_I)$ is a G_n/U_I-module, and one is led to ask if $E_n \wedge M_I$ is a G_n/U_I-spectrum. If so, then $E_n \wedge M_I$ is a discrete G_n-spectrum, and E_n is easily seen to be a continuous G_n-spectrum.

However, $E_n \wedge M_I$ is not a G_n/U-spectrum for all open normal subgroups U of G_n, as the following observation shows. As far as the author knows, this observation is due to Mike Hopkins; the author learned the proof from Hal Sadofsky.

Lemma 6.2. *There is no open normal subgroup U of G_n such that the G_n-action on $E_n \wedge M_I$ factors through G_n/U.*

Proof. Suppose the G_n-spectrum $E_n \wedge M_I$ is a G_n/U-spectrum. Then the G_n-action on the middle factor of $E_n \wedge E_n \wedge M_I$ factors through G_n/U, so that $\pi_*(E_n \wedge E_n \wedge M_I)$ is a G_n/U-module. Note that $\pi_*(E_n \wedge E_n \wedge M_I) \cong \mathrm{Map}_c^{\ell}(G_n, E_{n*}/I)$.

Since the G_n-module $\mathrm{Map}_c^{\ell}(G_n, E_{n*}/I)$ is a G_n/U-module, there is an isomorphism of sets $\mathrm{Map}_c^{\ell}(G_n, E_{n,0}/I) = \mathrm{Map}_c^{\ell}(G_n, E_{n,0}/I)^U \cong \mathrm{Map}_c(G_n/U, E_{n,0}/I)$. But the first set is infinite and the last is finite, a contradiction. $\square$

Since $\pi_*(E_n \wedge M_I)$ is a discrete G_n-module, one can still hope for a spectrum $E_n/I \simeq E_n \wedge M_I$, such that E_n/I is a discrete G_n-spectrum.

To produce E_n/I, we make the following observation. By [23, Remark 6.26], since U_i is an open normal subgroup of G_n, the presheaf $\mathrm{Hom}_{U_i}(-, (E_n/I)_{f,G_n})$ is fibrant in the model category of presheaves of spectra on the site $U_i\text{–}\mathbf{Sets}_{df}$. Thus, for each i, the map $E_n/I \to (E_n/I)_{f,G_n}$ is a trivial cofibration, with fibrant target, all in Sp_{U_i}, so that $((E_n/I)_{f,G_n})^{U_i} = (E_n/I)^{hU_i}$.

Combining this observation with the idea, discussed in §1, that $E_n \wedge M_I$ has homotopy fixed point spectra $(E_n \wedge M_I)^{hU_i} \simeq E_n^{\mathtt{h}U_i} \wedge M_I \simeq (E_n/I)^{hU_i}$, we have:

$$\begin{aligned} E_n/I &\simeq (E_n/I)_{f,G_n} \cong \operatorname{colim}_i ((E_n/I)_{f,G_n})^{U_i} = \operatorname{colim}_i (E_n/I)^{hU_i} \\ &\simeq \operatorname{colim}_i (E_n^{\mathtt{h}U_i} \wedge M_I) \cong F_n \wedge M_I. \end{aligned}$$

This argument suggests that $E_n \wedge M_I$ has the homotopy type of the discrete G_n-spectrum $F_n \wedge M_I$. To show that this is indeed the case, we consider the spectrum F_n in more detail. The key result is the following theorem, due to Devinatz and Hopkins.

Theorem 6.3 ([6])**.** *There is a weak equivalence $E_n \simeq \hat{L}(F_n)$.*

Proof. By [6, Thm. 3], $E_n^{h'\{e\}} \simeq E_n^{\mathtt{h}\{e\}}$. (We remark that this weak equivalence is far from obvious.) By [6, Definition 1.5], $E_n^{\mathtt{h}\{e\}} = \hat{L}(\operatorname{hocolim}_i E_n^{\mathtt{h}U_i})$, where the homotopy colimit is in the category E_∞. Then, by [6, Remark 1.6, Lemma 6.2], $\operatorname{hocolim}_i E_n^{\mathtt{h}U_i} \simeq \operatorname{colim}_i E_n^{\mathtt{h}U_i}$, where the colimit is in $\mathcal{M}_S$. Thus, as spectra in Sp, $E_n^{\mathtt{h}\{e\}} \simeq \hat{L}(F_n)$, so that $E_n \simeq E_n^{h'\{e\}} \simeq \hat{L}(F_n)$. $\square$

Corollary 6.4. *In the stable category, there are isomorphisms*

$$E_n \cong \operatorname{holim}_I (F_n \wedge M_I)_{\mathtt{f}} \cong \operatorname{holim}_I (E_n \wedge M_I)_{\mathtt{f}}.$$

The following result, a proof of which was first shown to me by Charles Rezk, shows that $E_n \wedge M_I \simeq F_n \wedge M_I$. This weak equivalence and $\mathrm{vcd}(G_n) < \infty$ are the main facts that make it possible to construct the homotopy fixed point spectra of E_n.

Corollary 6.5. *If Y is a finite spectrum of type n, then the G_n-equivariant map $F_n \wedge Y \to E_n \wedge Y$ is a weak equivalence. In particular, $E_n \wedge M_I \simeq F_n \wedge M_I$.*

Proof. We have $E_n \wedge Y \simeq \hat{L}(F_n) \wedge Y \simeq L_n(F_n) \wedge Y \simeq F_n \wedge Y$. □

Now we show that E_n is a continuous G_n-spectrum.

Theorem 6.6. *There is an isomorphism* $E_n \cong \operatorname{holim}_I (F_n \wedge M_I)_{f,G_n}$. *Thus,* E_n *is a continuous* G_n*-spectrum.*

Proof. By Corollary 6.4, $E_n \cong \operatorname{holim}_I (F_n \wedge M_I)_{\mathtt{f}}$. By functorial fibrant replacement, the map of towers $\{F_n \wedge M_I\} \to \{(F_n \wedge M_I)_{f,G_n}\}$ induces a map of towers $\{(F_n \wedge M_I)_{\mathtt{f}}\} \to \{((F_n \wedge M_I)_{f,G_n})_{\mathtt{f}}\}$ and, hence, weak equivalences

$$\operatorname{holim}_I (F_n \wedge M_I)_{f,G_n} \to \operatorname{holim}_I ((F_n \wedge M_I)_{f,G_n})_{\mathtt{f}} \leftarrow \operatorname{holim}_I (F_n \wedge M_I)_{\mathtt{f}}.$$

Thus, $\operatorname{holim}_I (F_n \wedge M_I)_{f,G_n}$ is isomorphic to $\operatorname{holim}_I (F_n \wedge M_I)_{\mathtt{f}}$ and E_n. Since $\{(F_n \wedge M_I)_{f,G_n}\}$ is a tower of fibrant spectra, $\operatorname{holim}_I (F_n \wedge M_I)_{f,G_n}$ is a continuous G_n-spectrum. Then, by Remark 4.6, E_n is a continuous G_n-spectrum. □

We conclude this section with some observations about F_n.

Lemma 6.7. *The map* $\eta\colon F_n \to E_n$ *is not a weak equivalence and* F_n *is not* $K(n)$*-local.*

Proof. If η is a weak equivalence, then $\pi_0(\eta)$ is a G_n-equivariant isomorphism from a discrete G_n-module (with all orbits finite) to a non-finite profinite G_n-module, which is impossible. If F_n is $K(n)$-local, then $F_n \simeq \hat{L}(F_n) \simeq E_n$, and η is a weak equivalence, a contradiction. □

Lemma 6.8. *The maps* $\hat{L}(F_n \wedge F_n) \to \hat{L}(E_n \wedge F_n) \to \hat{L}(E_n \wedge E_n)$ *are weak equivalences.*

Proof. Since $F_n \wedge M_I \simeq E_n \wedge M_I$, $F_n \wedge F_n \wedge M_I \simeq E_n \wedge E_n \wedge M_I$. Since $F_n \wedge F_n$, $E_n \wedge F_n$ and $E_n \wedge E_n$ are $E(n)$-local, the result follows from Theorem 2.8. □

Since $E_n^{\mathtt{h}U_i}$ is not an E_n-module (but E_n is an $E_n^{\mathtt{h}U_i}$-module), note that F_n and $\operatorname{holim}_I (F_n \wedge M_I)_{f,G_n}$ are not E_n-modules. However, any fully satisfactory model of E_n as a continuous G_n-spectrum would also be a twisted E_n-module G_n-spectrum, since E_n itself is such a spectrum. ("Twisted" means that the module structure map is G_n-equivariant, where $E_n \wedge E_n$ has the diagonal action.)

7. Homotopy fixed points when vcd(G) < ∞

In this section, G always has finite virtual cohomological dimension. Thus, there exists a uniform bound m, such that for all $K <_c G$, $\operatorname{vcd}(K) \leq m$. For $X \in Sp_G$, we use this fact to give a model for X^{hG} that eases the construction of its descent spectral sequence. Also, this fact yields a second model for X^{hK} that is functorial in K.

Definition 7.1. Consider the functor

$$\Gamma_G = \operatorname{Map}_c(G, -) \circ U\colon Sp_G \to Sp_G, \quad X \mapsto \Gamma_G(X) = \operatorname{Map}_c(G, X),$$

where $\Gamma_G(X)$ has the G-action defined in §3. We write Γ instead of Γ_G, when G is understood from context. There is a G-equivariant monomorphism $i\colon X \to \Gamma X$ defined, on the level of sets, by $i(x)(g) = g \cdot x$. As in [44, 8.6.2], since U and $\operatorname{Map}_c(G, -)$ are adjoints, Γ forms a triple and there is a cosimplicial discrete G-spectrum $\Gamma^\bullet X$, with $(\Gamma^\bullet X)^k \cong \operatorname{Map}_c(G^{k+1}, X)$.

We recall the construction of Thomason's hypercohomology spectrum for the topos **Shv** (see [41, 1.31-1.33] and [29, §1.3, §3.2] for more details). Consider the functor

$$T = u_*u^* \colon \mathbf{ShvSpt} \to \mathbf{ShvSpt}, \quad \mathcal{F} \mapsto \mathrm{Hom}_G(-, \mathrm{Map}_c(G, \operatorname{colim}_N \mathcal{F}(G/N))),$$

obtained by composing the adjoints in the point of the topos. Then, for $X \in Sp_G$, $T(\mathrm{Hom}_G(-,X)) \cong \mathrm{Hom}_G(-, \mathrm{Map}_c(G,X))$. By iterating this isomorphism, the cosimplicial sheaf of spectra $T^\bullet\mathrm{Hom}_G(-,X)$ gives rise to the cosimplicial sheaf $\mathrm{Hom}_G(-, \Gamma^\bullet X)$.

Definition 7.2. Given $X \in Sp_G$, the *presheaf of hypercohomology spectra of* $G-$**Sets**$_{df}$ *with coefficients in* X is the presheaf of spectra

$$\mathbb{H}^\bullet(-;X) = \operatorname{holim}_\Delta \mathrm{Hom}_G(-, \Gamma^\bullet X) \colon (G-\mathbf{Sets}_{df})^{\mathrm{op}} \to Sp,$$

and $\mathbb{H}^\bullet(S;X) = \operatorname{holim}_\Delta \mathrm{Hom}_G(S, \Gamma^\bullet X)$ is the hypercohomology spectrum of S with coefficients in X.

The map $X_f \to \Gamma^\bullet X_f$, induced by i, out of the constant cosimplicial diagram, and $\mathrm{Hom}_G(-,X_f) \to \lim_\Delta \mathrm{Hom}_G(-,X_f) \to \operatorname{holim}_\Delta \mathrm{Hom}_G(-,X_f)$ induce a canonical map $\mathrm{Hom}_G(-,X_f) \to \mathbb{H}^\bullet(-,X_f)$.

Now we show that $\mathbb{H}^\bullet(*,X_f)$ is a model for X^{hG}. Below, a *cosimplicial globally fibrant presheaf* is a cosimplicial presheaf of spectra that is globally fibrant at each level.

Lemma 7.3. *If* $\mathcal{F}^\bullet$ *is a cosimplicial globally fibrant presheaf, then* $\operatorname{holim}_\Delta \mathcal{F}^\bullet$ *is a globally fibrant presheaf.*

Proof. By [23, Rk. 2.35], this is equivalent to showing that, for each $n \geq 0$, (a) $\operatorname{holim}_\Delta (\mathcal{F}^\bullet)^n$ is a globally fibrant simplicial presheaf; and (b) the adjoint of the bonding map, the composition

$$\gamma \colon \operatorname{holim}_\Delta (\mathcal{F}^\bullet)^n \to \Omega(\operatorname{holim}_\Delta (\mathcal{F}^\bullet)^{n+1}) \cong \operatorname{holim}_\Delta \Omega(\mathcal{F}^\bullet)^{n+1}$$

is a local weak equivalence of simplicial presheaves. Part (a), the difficult part of this lemma, is proven in [21, Prop. 3.3].

The map γ is a local weak equivalence if the map of stalks $\operatorname{colim}_N \gamma(G/N)$ is a weak equivalence in $\mathcal{S}$, which is true if each $\gamma(G/N)$ is a weak equivalence. By [21, pg. 74], if P is a globally fibrant simplicial presheaf, then ΩP is too, so that $\sigma \colon (\mathcal{F}^\bullet)^n \to \Omega(\mathcal{F}^\bullet)^{n+1}$ is a cosimplicial diagram of local weak equivalences between globally fibrant simplicial presheaves. Thus, $\sigma(G/N)$ is a cosimplicial diagram of weak equivalences between Kan complexes, so that $\gamma(G/N) = \operatorname{holim}_\Delta \sigma(G/N)$ is indeed a weak equivalence. □

The following result is not original: it is basically a special case of [21, Prop. 3.3], versions of which appear in [11, §5], [29, Prop. 3.20], and [30, Prop. 6.1]. Since the result is central to our work, for the benefit of the reader, we give the details of the proof. For use now and later, we recall that, for any group G and any closed subgroup K, $H^s_c(K; \mathrm{Map}_c(G,A)) = 0$, when $s > 0$ and A is any discrete abelian group [45, Lemma 9.4.5].

Theorem 7.4. *Let* G *be a profinite group with* $\mathrm{vcd}(G) \leq m$, *and let* X *be a discrete* G*-spectrum. Then there are weak equivalences*

$$\mathrm{Hom}_G(-,X) \xrightarrow{\simeq} \mathrm{Hom}_G(-,X_f) \xrightarrow{\simeq} \operatorname{holim}_\Delta \mathrm{Hom}_G(-, \Gamma^\bullet X_f),$$

and $\operatorname{holim}_\Delta \operatorname{Hom}_G(-,\Gamma^\bullet X_f)$ *is a globally fibrant model for* $\operatorname{Hom}_G(-,X)$*. Thus, evaluation at* $* \in G-\mathbf{Sets}_{df}$ *gives a weak equivalence* $X^{hG} \to \operatorname{holim}_\Delta (\Gamma^\bullet X_f)^G$*.*

Remark 7.5. The weak equivalence $X \cong \operatorname{colim}_N X^N \to \operatorname{colim}_N (X^N)_{\mathtt{f}}$, whose target is a discrete G-spectrum that is fibrant in Sp, induces a weak equivalence $X_{f,G} \to (\operatorname{colim}_N (X^N)_{\mathtt{f}})_{f,G}$. Thus, there are weak equivalences

$$\mathbb{H}^\bullet(*, X_f) \to \mathbb{H}^\bullet(*, (\operatorname{colim}_N (X^N)_{\mathtt{f}})_{f,G}) \leftarrow \mathbb{H}^\bullet(*, \operatorname{colim}_N (X^N)_{\mathtt{f}}),$$

so that $\mathbb{H}^\bullet(*, \operatorname{colim}_N (X^N)_{\mathtt{f}})$ is a model for X^{hG} that does not require the model category Sp_G for its construction.

Proof of Theorem 7.4. Since X_f is fibrant in Sp, ΓX_f is fibrant in Sp_G by Corollary 3.7. By iteration, $\operatorname{Hom}_G(-,\Gamma^\bullet X_f)$ is a cosimplicial globally fibrant presheaf, so that $\operatorname{holim}_\Delta \operatorname{Hom}_G(-,\Gamma^\bullet X_f)$ is globally fibrant, by Lemma 7.3. It only remains to show that $\lambda\colon \operatorname{Hom}_G(-,X) \to \operatorname{holim}_\Delta \operatorname{Hom}_G(-,\Gamma^\bullet X_f)$ is a weak equivalence.

By hypothesis, G contains an open subgroup H with $\operatorname{cd}(H) \le m$. Then by [45, Lem. 0.3.2], H contains a subgroup K that is an open normal subgroup of G. Let $\{N\}$ be the collection of open normal subgroups of G. Let $N' = N \cap K$. Observe that $\{N'\}$ is a cofinal subcollection of open normal subgroups of G so that $G \cong \lim_{N'} G/N'$. Since $N' <_c H$, $\operatorname{cd}(N') \le \operatorname{cd}(H)$. Thus, $H^s_c(N';M) = 0$, for all $s > m+1$, whenever M is a discrete N'-module. Henceforth, we drop the $'$ from N' to ease the notation: N is really $N \cap K$.

Any presheaf of sets $\mathcal{F}$ has stalk $\operatorname{colim}_N \mathcal{F}(G/N)$, so that λ is a weak equivalence if $\lambda_u\colon X \cong \operatorname{colim}_N X^N \to \operatorname{colim}_N \operatorname{holim}_\Delta (\Gamma^\bullet X_f)^N$ is a weak equivalence. Since $\operatorname{Hom}_G(-,\Gamma^\bullet X_f)$ is a cosimplicial globally fibrant spectrum, the diagram $(\Gamma^\bullet X_f)^N$ is a cosimplicial fibrant spectrum. Then, for each N, there is a conditionally convergent spectral sequence

$$E_2^{s,t}(N) = \pi^s\pi_t((\Gamma^\bullet X_f)^N) \Rightarrow \pi_{t-s}(\operatorname{holim}_\Delta (\Gamma^\bullet X_f)^N). \tag{7.6}$$

Because $\pi_t(X)$ is a discrete G-module, we have

$$\pi_t(\operatorname{Map}_c(G,X_f)^N) \cong \pi_t(\operatorname{Map}_c(G/N,X_f)) \cong \textstyle\prod_{G/N} \pi_t(X) \cong \operatorname{Map}_c(G,\pi_t(X))^N$$

and

$$\pi_t(\operatorname{Map}_c(G,X_f)) \cong \pi_t(\operatorname{colim}_N \textstyle\prod_{G/N} X_f) \cong \operatorname{Map}_c(G,\pi_t(X)).$$

Iterating such manipulations, we obtain $\pi^s\pi_t((\Gamma^\bullet X_f)^N) \cong H^s((\Gamma^*\pi_t(X))^N)$. The cochain complex $0 \to \pi_t(X) \to \Gamma^*\pi_t(X)$ of discrete N-modules is exact (see e.g. [11, pp. 210-211]), and, for $k \ge 1$ and $s > 0$,

$$H^s_c(N;\Gamma^k\pi_t(X)) \cong H^s_c(N;\operatorname{Map}_c(G,\Gamma^{k-1}\pi_t(X))) = 0.$$

Thus, the above cochain complex is a resolution of $\pi_t(X)$ by $(-)^N$-acyclic modules, so that $E_2^{s,t}(N) \cong H^s_c(N;\pi_t(X))$. Taking a colimit over $\{N\}$ of (7.6) gives the spectral sequence

$$E_2^{s,t} = \operatorname{colim}_N H^s_c(N;\pi_t(X)) \Rightarrow \pi_{t-s}(\operatorname{colim}_N \operatorname{holim}_\Delta (\Gamma^\bullet X_f)^N). \tag{7.7}$$

Since $E_2^{s,*}(N) = 0$ whenever $s > m+1$, the E_2-terms $E_2(N)$ are uniformly bounded on the right. Therefore, by [29, Prop. 3.3], the colimit of the spectral sequences does converge to the colimit of the abutments, as asserted in (7.7).

Finally, $E_2^{*,t} \cong H^*_c(\lim_N N;\pi_t(X)) \cong H^*(\{e\};\pi_t(X))$, which is isomorphic to $\pi_t(X)$, concentrated in degree zero. Thus, (7.7) collapses and for all t,

$$\pi_t(\operatorname{colim}_N \operatorname{holim}_\Delta (\Gamma^\bullet X_f)^N) \cong \pi_t(X),$$

and hence, λ_u is a weak equivalence. □

Remark 7.8. Because of Theorem 7.4, if $\mathrm{vcd}(G) < \infty$ and X is a discrete G-spectrum, we make the identification

$$X^{hG} = \mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_G X_{f,G})^G = \mathbb{H}^{\bullet}(*, X_{f,G}).$$

In [43, §2.14], an expression that is basically equivalent to $\mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_G X_{f,G})^G$ is defined to be the homotopy fixed point spectrum X^{hG}, even if $\mathrm{vcd}(G) = \infty$. This approach has the disadvantage that $(-)^{hG}$ need not always be the total right derived functor of $(-)^G$. Thus, we only make the identification of Remark 7.8 when $\mathrm{vcd}(G) < \infty$.

Now it is easy to construct the descent spectral sequence. Note that if X is a discrete G-spectrum, the proof of Theorem 7.4 shows that

$$\pi^s \pi_t((\Gamma^{\bullet}_G X)^G) \cong \pi^s((\Gamma^{\bullet}_G \pi_t(X))^G) \cong H^s_c(G; \pi_t(X)).$$

Theorem 7.9. *If* $\mathrm{vcd}(G) < \infty$ *and* X *is a discrete* G*-spectrum, then there is a conditionally convergent descent spectral sequence*

$$E_2^{s,t} = H^s_c(G; \pi_t(X)) \Rightarrow \pi_{t-s}(X^{hG}).$$

Proof. As in Theorem 7.4, $(\Gamma^{\bullet} X_f)^G$ is a cosimplicial fibrant spectrum. Thus, we can form the homotopy spectral sequence for $\pi_*(\mathrm{holim}_{\Delta}\, (\Gamma^{\bullet} X_f)^G)$. □

Remark 7.10. The descent spectral sequence of Theorem 7.9 has been constructed in other contexts: for simplicial presheaves, presheaves of spectra, and S_G, see [21, Cor. 3.6], [23, §6.1], and [11, §§4, 5], respectively. In several of these examples, a Postnikov tower provides an alternative to the hypercohomology spectrum that we use. In all of these constructions of the descent spectral sequence, some kind of finiteness assumption is required in order to identify the homotopy groups of the abutment as being those of the homotopy fixed point spectrum.

Let X be a discrete G-spectrum. We now develop a second model for X^{hK}, where K is a closed subgroup of G, that is functorial in K.

The map $X \to X_{f,G}$ in Sp_K gives a weak equivalence $X^{hK} \to (X_{f,G})^{hK}$. Composition with the weak equivalence $(X_{f,G})^{hK} \to \mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_K((X_{f,G})_{f,K}))^K$ gives a weak equivalence $X^{hK} \to \mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_K((X_{f,G})_{f,K}))^K$ between fibrant spectra. The inclusion $K \to G$ induces a morphism $\Gamma_G(X_{f,G}) \to \Gamma_K(X_{f,G})$, giving a map $\Gamma^{\bullet}_G(X_{f,G}) \to \Gamma^{\bullet}_K(X_{f,G})$ of cosimplicial discrete K-spectra.

Lemma 7.11. *There is a weak equivalence*

$$\rho\colon \mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_G(X_{f,G}))^K \to \mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_K(X_{f,G}))^K \to \mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_K((X_{f,G})_{f,K}))^K.$$

Proof. Recall the conditionally convergent spectral sequence

$$H^s_c(K; \pi_t(X)) \cong H^s_c(K; \pi_t((X_{f,G})_{f,K})) \Rightarrow \pi_{t-s}(\mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_K((X_{f,G})_{f,K}))^K).$$

We compare this spectral sequence with the homotopy spectral sequence for $\mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_G(X_{f,G}))^K$. Note that if Y is a discrete G-spectrum that is fibrant as a spectrum, then $\mathrm{Map}_c(G, Y) \cong \mathrm{colim}_N \prod_{G/N} Y$ and

$$\mathrm{Map}_c(G, Y)^K \cong \mathrm{Map}_c(G/K, Y) \cong \mathrm{colim}_N \prod_{G/(NK)} Y$$

are fibrant spectra. Thus, $(\Gamma^{\bullet}_G(X_{f,G}))^K$ is a cosimplicial fibrant spectrum, and there is a conditionally convergent spectral sequence

$$E_2^{s,t} = H^s((\Gamma^{*}_G \pi_t(X_{f,G}))^K) \Rightarrow \pi_{t-s}(\mathrm{holim}_{\Delta}\, (\Gamma^{\bullet}_G(X_{f,G}))^K).$$

As in the proof of Theorem 7.4, $0 \to \pi_t(X_{f,G}) \to \Gamma^*_G(\pi_t(X_{f,G}))$ is a $(-)^K$-acyclic resolution of $\pi_t(X_{f,G})$. Hence,

$$E_2^{s,t} \cong H^s_c(K; \pi_t(X_{f,G})) \cong H^s_c(K; \pi_t(X)).$$

Since ρ is compatible with the isomorphism between the two E_2-terms, the spectral sequences are isomorphic and ρ is a weak equivalence. □

Remark 7.12. Lemma 7.11 gives the following weak equivalences between fibrant spectra:

$$X^{hK} = (X_{f,K})^K \to \operatorname{holim}_\Delta (\Gamma^\bullet_K((X_{f,G})_{f,K}))^K \leftarrow \operatorname{holim}_\Delta (\Gamma^\bullet_G(X_{f,G}))^K.$$

Thus, if $\mathrm{vcd}(G) < \infty$, X is a discrete G-spectrum, and K is a closed subgroup of G, then $\operatorname{holim}_\Delta (\Gamma^\bullet_G(X_{f,G}))^K$ is a model for X^{hK}, so that

$$X^{hK} = \operatorname{holim}_\Delta (\Gamma^\bullet_G(X_{f,G}))^K$$

is another definition of the homotopy fixed points.

This discussion yields the following result.

Theorem 7.13. *If X is a discrete G-spectrum, with* $\mathrm{vcd}(G) < \infty$*, then there is a presheaf of spectra $P(X)\colon (\mathcal{O}_G)^{\mathrm{op}} \to Sp$, defined by*

$$P(X)(G/K) = \operatorname{holim}_\Delta (\Gamma^\bullet_G(X_{f,G}))^K = X^{hK}.$$

Proof. If Y is a discrete G-set, any morphism $f\colon G/H \to G/K$, in $\mathcal{O}_G$, induces a map

$$\operatorname{Map}_c(G,Y)^K \cong \operatorname{Map}_c(G/K,Y) \to \operatorname{Map}_c(G/H,Y) \cong \operatorname{Map}_c(G,Y)^H.$$

Thus, if $Y \in Sp_G$, f induces a map $\operatorname{Map}_c(G,Y)^K \to \operatorname{Map}_c(G,Y)^H$, so that there is a map

$$P(X)(f)\colon \operatorname{holim}_\Delta (\Gamma^\bullet_G(X_{f,G}))^K \to \operatorname{holim}_\Delta (\Gamma^\bullet_G(X_{f,G}))^H.$$

It is easy to check that $P(X)$ is actually a functor. □

We conclude this section by pointing out a useful fact: smashing with a finite spectrum, with trivial G-action, commutes with taking homotopy fixed points. To state this precisely, we first of all define the relevant map.

Let X be a discrete G-spectrum and let Y be any spectrum with trivial G-action. Then there is a map

$$(\operatorname{holim}_\Delta (\Gamma^\bullet_G X_f)^G) \wedge Y \to \operatorname{holim}_\Delta ((\Gamma^\bullet_G X_f)^G \wedge Y) \to \operatorname{holim}_\Delta ((\Gamma^\bullet_G X_f) \wedge Y)^G.$$

Also, there is a natural G-equivariant map $\operatorname{Map}_c(G,X) \wedge Y \to \operatorname{Map}_c(G, X \wedge Y)$ that is defined by the composition

$$(\operatorname{colim}_N \textstyle\prod_{G/N} X) \wedge Y \cong \operatorname{colim}_N ((\prod_{G/N} X) \wedge Y) \to \operatorname{colim}_N \prod_{G/N} (X \wedge Y),$$

by using the isomorphism $\operatorname{Map}_c(G,X) \cong \operatorname{colim}_N \prod_{G/N} X$. This gives a natural G-equivariant map

$$(\Gamma_G \Gamma_G X) \wedge Y \to \Gamma_G ((\Gamma_G X) \wedge Y) \to \Gamma_G \Gamma_G (X \wedge Y).$$

Thus, iteration gives a G-equivariant map of cosimplicial spectra

$$(\Gamma^\bullet X) \wedge Y \to \Gamma^\bullet (X \wedge Y).$$

Therefore, if $\mathrm{vcd}(G) < \infty$, there is a canonical map $X^{hG} \wedge Y \to (X_f \wedge Y)^{hG}$ that is defined by composing the map $X^{hG} \wedge Y \to \mathrm{holim}_{\Delta}\, ((\Gamma^\bullet X_f) \wedge Y)^G$, from above, with the map

$$\mathrm{holim}_{\Delta}\, ((\Gamma^\bullet X_f) \wedge Y)^G \to \mathrm{holim}_{\Delta}\, (\Gamma^\bullet(X_f \wedge Y))^G \to \mathrm{holim}_{\Delta}\, (\Gamma^\bullet(X_f \wedge Y)_f)^G.$$

Lemma 7.14 ([29, Prop. 3.10])**.** *If $\mathrm{vcd}(G) < \infty$, $X \in Sp_G$, and Y is a finite spectrum with trivial G-action, then $X^{hG} \wedge Y \to (X_{f,G} \wedge Y)^{hG}$ is a weak equivalence.*

Remark 7.15. By Lemma 7.14, when Y is a finite spectrum, there is a zigzag of natural weak equivalences $X^{hG} \wedge Y \to (X_{f,G} \wedge Y)^{hG} \leftarrow (X \wedge Y)^{hG}$. We refer to this zigzag by writing $X^{hG} \wedge Y \cong (X \wedge Y)^{hG}$.

8. Homotopy fixed points for towers in Sp_G

In this section, $\{Z_i\}$ is always a tower in Sp_G (except in Definition 8.9). For $\{Z_i\}$ a tower of fibrant spectra, we define the homotopy fixed point spectrum $(\mathrm{holim}_i\, Z_i)^{hG}$ and construct its descent spectral sequence.

Definition 8.1. If $\{Z_i\}$ in $\mathbf{tow}(Sp_G)$ is a tower of fibrant spectra, we define $Z = \mathrm{holim}_i\, Z_i$, a continuous G-spectrum. The homotopy fixed point spectrum Z^{hG} is defined to be $\mathrm{holim}_i\, Z_i^{hG}$, a fibrant spectrum.

We make several comments about Definition 8.1. Let H be a closed subgroup of G. Then the maps

$$\mathrm{holim}_i\, ((Z_i)_f)^H \to \mathrm{holim}_i\, \mathrm{holim}_{\Delta}\, (\Gamma^\bullet_H (Z_i)_f)^H \quad \text{and}$$

$$\mathrm{holim}_i\, \mathrm{holim}_{\Delta}\, (\Gamma^\bullet_G (Z_i)_{f,G})^H \to \mathrm{holim}_i\, \mathrm{holim}_{\Delta}\, (\Gamma^\bullet_H ((Z_i)_{f,G})_{f,H})^H$$

are weak equivalences. Thus, in Definition 8.1, any one of our three definitions for homotopy fixed points (Definition 5.2, Remarks 7.8 and 7.12) can be used for Z_i^{hH}.

In Definition 8.1, suppose that not all the Z_i are fibrant in Sp. Then the map $Z = \mathrm{holim}_i\, Z_i \to \mathrm{holim}_i\, (Z_i)_{f,\{e\}} = \mathrm{holim}_i\, Z_i^{h\{e\}} = Z^{h\{e\}}$ need not be a weak equivalence. Thus, for an arbitrary tower in Sp_G, Definition 8.1 can fail to have the desired property that $Z \to Z^{h\{e\}}$ is a weak equivalence.

Below, Lemmas 8.2 and 8.3, and Remark 8.4, show that when G is a finite group, $Z^{hG} \simeq Z^{h'G}$, and, for any G, Z^{hG} can be obtained by using a total right derived functor that comes from fixed points. Thus, Definition 8.1 generalizes the notion of homotopy fixed points to towers of discrete G-spectra.

Lemma 8.2. *Let G be a finite group and let $\{Z_i\}$ in $\mathbf{tow}(Sp_G)$ be a tower of fibrant spectra. Then there is a weak equivalence $Z^{hG} \to Z^{h'G}$.*

Proof. We have: $Z^{hG} = \mathrm{holim}_i\, \lim_G\, (Z_i)_f$ and $Z^{h'G} = \mathrm{holim}_G\, \mathrm{holim}_i\, (Z_i)_f$ (since $Z = \mathrm{holim}_i\, Z_i \to \mathrm{holim}_i\, (Z_i)_f$ is a weak equivalence and G-equivariant, and $\mathrm{holim}_i\, (Z_i)_f$ is fibrant in Sp). Then the map $Z^{hG} \to Z^{h'G}$ is defined to be

$$\mathrm{holim}_i\, \lim_G\, (Z_i)_f \to \mathrm{holim}_i\, \mathrm{holim}_G\, (Z_i)_f \cong \mathrm{holim}_G\, \mathrm{holim}_i\, (Z_i)_f.$$

For each i, $\lim_G\, (Z_i)_f \to \mathrm{holim}_G\, (Z_i)_f$ is a weak equivalence between fibrant spectra, so that the desired map is a weak equivalence. □

In the lemma below,

$$\mathbf{R}(\lim_i\, (-)^G)\colon \mathrm{Ho}(\mathbf{tow}(Sp_G)) \to \mathrm{Ho}(Sp)$$

is the total right derived functor of the functor $\lim_i\, (-)^G\colon \mathbf{tow}(Sp_G) \to Sp$.

Lemma 8.3. *If $\{Z_i\}$ is an arbitrary tower in Sp_G, then*

$$\operatorname{holim}_i ((Z_i)_f)^G \xrightarrow{\simeq} \operatorname{holim}_i ((Z_i)'_f)^G \xleftarrow{\simeq} \lim_i ((Z_i)'_f)^G = \mathbf{R}(\lim_i (-)^G)(\{Z_i\}).$$

Proof. The first weak equivalence follows from $((Z_i)_f)^G \to ((Z_i)'_f)^G$ being a weak equivalence between fibrant spectra. The second weak equivalence holds because $\{((Z_i)'_f)^G\}$ is a tower of fibrant spectra, such that all maps in the tower are fibrations. Finally, the equality is because $\{Z_i\} \to \{(Z_i)'_f\}$ is a trivial cofibration in $\mathbf{tow}(Sp_G)$. □

Remark 8.4. By Lemma 5.3, if $X \in Sp_G$, then $X^{hG} = (\mathbf{R}(-)^G)(X)$. Also, by Lemma 8.3, if $\{Z_i\}$ in $\mathbf{tow}(Sp_G)$ is a tower of fibrant spectra, then

$$Z^{hG} = \operatorname{holim}_i Z_i^{hG} = \operatorname{holim}_i ((Z_i)_f)^G \cong \mathbf{R}(\lim_i (-)^G)(\{Z_i\}).$$

Thus, the homotopy fixed point spectrum Z^{hG} is again given by the total right derived functor of an appropriately defined functor involving G-fixed points.

Given any tower in Sp_G of fibrant spectra, there is a descent spectral sequence whose E_2-term is a version of continuous cohomology, whose definition we now recall from [20]. We use $\mathrm{DMod}(G)$ to denote the category of discrete G-modules.

Definition 8.5. Let $\mathrm{DMod}(G)^{\mathbb{N}}$ denote the category of diagrams in discrete G-modules of the form $\cdots \to M_2 \to M_1 \to M_0$. Then $H^s_{\mathrm{cont}}(G; \{M_i\})$, the continuous cohomology of G with coefficients in the tower $\{M_i\}$, is the sth right derived functor of the left exact functor

$$\lim_i (-)^G \colon \mathrm{DMod}(G)^{\mathbb{N}} \to \mathbf{Ab}, \quad \{M_i\} \mapsto \lim_i M_i^G.$$

By [20, Theorem 2.2], if the tower of abelian groups $\{M_i\}$ satisfies the Mittag-Leffler condition, then $H^s_{\mathrm{cont}}(G; \{M_i\}) \cong H^s_{\mathrm{cts}}(G; \lim_i M_i)$, for $s \geq 0$.

Theorem 8.6. *If $\mathrm{vcd}(G) < \infty$ and $\{Z_i\}$ in $\mathbf{tow}(Sp_G)$ is a tower of fibrant spectra, then there is a conditionally convergent descent spectral sequence*

$$H^s_{\mathrm{cont}}(G; \{\pi_t(Z_i)\}) \Rightarrow \pi_{t-s}(Z^{hG}). \tag{8.7}$$

Remark 8.8. Such a spectral sequence goes back to Thomason's construction of the ℓ-adic descent spectral sequence of algebraic K-theory ([41], [29]). Spectral sequence (8.7) is the homotopy spectral sequence

$$E_2^{s,t} = \lim{}^s_{\Delta \times \{i\}} \pi_t((\Gamma^\bullet_G((Z_i)_{f,G}))^G) \Rightarrow \pi_{t-s}(\operatorname{holim}_{\Delta \times \{i\}} (\Gamma^\bullet_G((Z_i)_{f,G}))^G),$$

where the identification $E_2^{s,t} \cong H^s_{\mathrm{cont}}(G; \{\pi_t(Z_i)\})$ depends on the fact that if $\{I_i\}$ is an injective object in $\mathrm{DMod}(G)^{\mathbb{N}}$, then $\lim{}^s_{\Delta \times \{i\}} (\Gamma^\bullet_G I_i)^G = 0$, for $s > 0$. We omit the details of the proof, since it is a special case of [9, Prop. 3.1.2], and also because (8.7) is not our focus of interest.

For our applications, instead of spectral sequence (8.7), we are more interested in descent spectral sequence (8.11) below. Spectral sequence (8.11), a homotopy spectral sequence for a particular cosimplicial spectrum, is more suitable for comparison with the $K(n)$-local E_n-Adams spectral sequence (see [6, Prop. A.5]), when (8.11) has abutment $\pi_*((E_n \wedge X)^{hG})$, where X is a finite spectrum.

Definition 8.9. If $\{Z_i\}$ is a tower of spectra such that $\{\pi_t(Z_i)\}$ satisfies the Mittag-Leffler condition for every $t \in \mathbb{Z}$, then $\{Z_i\}$ is a *Mittag-Leffler tower* of spectra.

Theorem 8.10. *If* $\mathrm{vcd}(G) < \infty$ *and* $\{Z_i\}$ *in* $\mathbf{tow}(Sp_G)$ *is a tower of fibrant spectra, then there is a conditionally convergent descent spectral sequence*

$$E_2^{s,t} = \pi^s\pi_t(\mathrm{holim}_i\,(\Gamma^\bullet_G(Z_i)_f)^G) \Rightarrow \pi_{t-s}(Z^{hG}). \tag{8.11}$$

If $\{Z_i\}$ *is a Mittag-Leffler tower, then* $E_2^{s,t} \cong H^s_{\mathrm{cts}}(G; \pi_t(Z))$.

Remark 8.12. In Theorem 8.10, when $\{Z_i\}$ is a Mittag-Leffler tower, $E_2^{s,t} \cong H^s_{\mathrm{cts}}(G;\pi_t(Z)) \cong H^s_{\mathrm{cont}}(G;\{\pi_t(Z_i)\})$, and spectral sequence (8.11) is identical to (8.7). However, in general, spectral sequences (8.7) and (8.11) are different. For example, if $G = \{e\}$, then in (8.7), $E_2^{0,t} = \lim_i \pi_t(Z_i)$, whereas in (8.11), $E_2^{0,t} = \pi_t(\mathrm{holim}_i\, Z_i)$.

Proof of Theorem 8.10. Note that $Z^{hG} \cong \mathrm{holim}_\Delta\, \mathrm{holim}_i\,(\Gamma^\bullet_G(Z_i)_f)^G$, and the diagram $\mathrm{holim}_i\,(\Gamma^\bullet_G(Z_i)_f)^G$ is a cosimplicial fibrant spectrum.

Let $\{Z_i\}$ be a Mittag-Leffler tower. For $k \geq 0$, Lemma 2.10 and Remark 2.11 imply that $\lim^1{}_i\, \mathrm{Map}_c(G^k, \pi_{t+1}(Z_i)) = 0$. Therefore,

$$\pi_t(\mathrm{holim}_i\,(\mathrm{Map}_c(G^{k+1},(Z_i)_f))^G) \cong \lim_i \mathrm{Map}_c(G^{k+1},\pi_t(Z_i))^G,$$

and hence, $\pi_t(\mathrm{holim}_i\,(\Gamma^\bullet(Z_i)_f)^G) \cong \lim_i\,(\Gamma^\bullet\pi_t(Z_i))^G$. Thus,

$$E_2^{s,t} \cong \pi^s(\lim_i\,(\Gamma^\bullet\pi_t(Z_i))^G) \cong H^s(\lim_i\,(-)^G\,\{\Gamma^*\pi_t(Z_i)\}_i).$$

Consider the exact sequence $\{0\} \to \{\pi_t(Z_i)\} \to \{\Gamma^*\pi_t(Z_i)\}$ in $\mathrm{DMod}(G)^{\mathbb{N}}$. Note that, for $s, k > 0$, by Theorem 2.15,

$$H^s_{\mathrm{cont}}(G;\{\Gamma^k\pi_t(Z_i)\}) \cong H^s_{\mathrm{cts}}(G;\lim_i \Gamma^k\pi_t(Z_i)) \cong \lim_i H^s_c(G;\Gamma^k\pi_t(Z_i)) = 0,$$

since the tower $\{\Gamma^k\pi_t(Z_i)\}$ satisfies the Mittag-Leffler condition, and, for each i, $\Gamma^k\pi_t(Z_i) \cong \mathrm{Map}_c(G,\Gamma^{k-1}\pi_t(Z_i))$ is $(-)^G$-acyclic. Thus, the above exact sequence is a $(\lim_i\,(-)^G)$-acyclic resolution of $\{\pi_t(Z_i)\}$, so that we obtain the isomorphism $E_2^{s,t} \cong H^s_{\mathrm{cont}}(G;\{\pi_t(Z_i)\}) \cong H^s_{\mathrm{cts}}(G;\pi_t(Z))$. □

By Remark 8.4, we can rewrite spectral sequence (8.11), when $\{Z_i\}$ is a Mittag-Leffler tower, in a more conceptual way:

$$R^s(\lim_i\,(-)^G)\{\pi_t(Z_i)\} \Rightarrow \pi_{t-s}(\mathbf{R}(\lim_i\,(-)^G)(\{Z_i\})).$$

Spectral sequence (8.7) can always be written in this way.

9. HOMOTOPY FIXED POINT SPECTRA FOR $\hat{L}(E_n \wedge X)$

In this section, for any spectrum X and any closed subgroup G of G_n, we define the homotopy fixed point spectrum $(\hat{L}(E_n \wedge X))^{hG}$, using the continuous action of G.

Let X be an arbitrary spectrum with trivial G_n-action. By Corollary 6.5, there is a weak equivalence $F_n \wedge M_I \wedge X \to E_n \wedge M_I \wedge X$. Then, using functorial fibrant replacement, there is a map of towers

$$\{(F_n \wedge M_I \wedge X)_{\mathtt{f}}\} \to \{(E_n \wedge M_I \wedge X)_{\mathtt{f}}\},$$

which yields the weak equivalence

$$\hat{L}(E_n \wedge X) \cong \mathrm{holim}_I\,(E_n \wedge M_I \wedge X)_{\mathtt{f}} \xleftarrow{\simeq} \mathrm{holim}_I\,(F_n \wedge M_I \wedge X)_{\mathtt{f}}.$$

As in the proof of Theorem 6.6, this implies the following lemma, since the diagram $\{(F_n \wedge M_I \wedge X)_{f,G_n}\}$ is a tower of fibrant spectra.

Lemma 9.1. *Given any spectrum X with trivial G_n-action, the isomorphism*

$$\hat{L}(E_n \wedge X) \cong \operatorname{holim}_I (F_n \wedge M_I \wedge X)_{f,G_n}$$

makes $\hat{L}(E_n \wedge X)$ a continuous G_n-spectrum.

Let G be any closed subgroup of G_n. Since $\{(F_n \wedge M_I \wedge X)_{f,G_n}\}$ is a tower of discrete G-spectra that are fibrant in Sp, Lemma 9.1 also shows that $\hat{L}(E_n \wedge X)$ is a continuous G-spectrum. By Theorem 3.6, the composition

$$(F_n \wedge M_I \wedge X) \to (F_n \wedge M_I \wedge X)_{f,G_n} \to ((F_n \wedge M_I \wedge X)_{f,G_n})_{f,G}$$

is a trivial cofibration in Sp_G, with target fibrant in Sp_G. Therefore,

$$((F_n \wedge M_I \wedge X)_{f,G_n})^{hG} = (F_n \wedge M_I \wedge X)^{hG}.$$

Then, by Definition 8.1, we have the following.

Definition 9.2. Let $G <_c G_n$. Then

$$E_n^{hG} = (\operatorname{holim}_I (F_n \wedge M_I)_{f,G_n})^{hG} = \operatorname{holim}_I (F_n \wedge M_I)^{hG}.$$

More generally, for any spectrum X,

$$(\hat{L}(E_n \wedge X))^{hG} = (\operatorname{holim}_I (F_n \wedge M_I \wedge X)_{f,G_n})^{hG} = \operatorname{holim}_I (F_n \wedge M_I \wedge X)^{hG}.$$

Remark 9.3. When X is a finite spectrum, $E_n \wedge X \simeq \hat{L}(E_n \wedge X)$. Thus, we have $(E_n \wedge X)^{hG} \cong (\hat{L}(E_n \wedge X))^{hG}$.

Remark 9.4. For any X, $\hat{L}(E_n \wedge X) \cong \operatorname{holim}_I (F_n \wedge M_I \wedge X)_{f,G}$ also shows that $\hat{L}(E_n \wedge X)$ is a continuous G-spectrum. Note that, by definition, the spectra $((F_n \wedge M_I \wedge X)_{f,G})^{hG}$ and $(F_n \wedge M_I \wedge X)^{hG}$ are identical. This implies that

$$(\hat{L}(E_n \wedge X))^{hG} = \operatorname{holim}_I ((F_n \wedge M_I \wedge X)_{f,G})^{hG} = \operatorname{holim}_I (F_n \wedge M_I \wedge X)^{hG},$$

as before.

Note that Definition 9.2 implies the identifications

$$E_n^{hG} = \operatorname{holim}_I \operatorname{holim}_\Delta (\Gamma^\bullet_{G_n}(F_n \wedge M_I)_{f,G_n})^G$$

and

$$(\hat{L}(E_n \wedge X))^{hG} = \operatorname{holim}_I \operatorname{holim}_\Delta (\Gamma^\bullet_{G_n}(F_n \wedge M_I \wedge X)_{f,G_n})^G.$$

The first identification, coupled with Theorem 7.13, implies the following result.

Theorem 9.5. *There is a functor*

$$P\colon (\mathcal{O}_{G_n})^{\mathrm{op}} \to Sp, \quad P(G_n/G) = E_n^{hG},$$

where G is any closed subgroup of G_n.

In addition to the above identifications, we also have

$$E_n^{hG} = \operatorname{holim}_I \operatorname{holim}_\Delta (\Gamma^\bullet_G(F_n \wedge M_I)_{f,G})^G$$

and

$$(\hat{L}(E_n \wedge X))^{hG} = \operatorname{holim}_I \operatorname{holim}_\Delta (\Gamma^\bullet_G(F_n \wedge M_I \wedge X)_{f,G})^G.$$

Since $E_n^{\mathrm{h}G}$ is $K(n)$-local, one would expect that E_n^{hG} is $K(n)$-local; this is verified below.

Lemma 9.6. *Let G be a closed subgroup of G_n and let X be any spectrum. Then the spectra $(F_n \wedge M_I \wedge X)^{hG}$ and $(\hat{L}(E_n \wedge X))^{hG}$ are $K(n)$-local. Also, $(F_n \wedge X)^{hG}$ is $E(n)$-local.*

Proof. Recall that $(F_n \wedge M_I \wedge X)^{hG} = \operatorname{holim}_\Delta (\Gamma^\bullet_G (F_n \wedge M_I \wedge X)_{f,G})^G$, and, for $k \geq 1$,

$$(\Gamma^k_G(F_n \wedge M_I \wedge X)_f)^G \simeq \operatorname{Map}_c(G^{k-1}, F_n \wedge X) \wedge M_I.$$

Since $\operatorname{Map}_c(G, F_n \wedge X) \cong \operatorname{colim}_i \prod_{G/(U_i \cap G)} (F_n \wedge X)$, and $F_n \wedge X$ is $E(n)$-local, the finite product is too, and hence, the direct limit $\operatorname{Map}_c(G, F_n \wedge X)$ is $E(n)$-local. Iterating this argument shows that $\operatorname{Map}_c(G^{k-1}, F_n \wedge X) \cong \Gamma_G \Gamma_G \cdots \Gamma_G(F_n \wedge X)$ is $E(n)$-local. Smashing $\operatorname{Map}_c(G^{k-1}, F_n \wedge X)$ with the type n spectrum M_I shows that $(\Gamma^k_G(F_n \wedge M_I \wedge X)_f)^G$ is $K(n)$-local. Therefore, since the homotopy limit of an arbitrary diagram of E-local spectra is E-local [1, pg. 259], $(F_n \wedge M_I \wedge X)^{hG}$ and $(\hat{L}(E_n \wedge X))^{hG}$ are $K(n)$-local. The same argument shows that $(F_n \wedge X)^{hG}$ is $E(n)$-local. □

The following theorem shows that the homotopy fixed points of $\hat{L}(E_n \wedge X)$ are obtained by taking the $K(n)$-localization of the homotopy fixed points of the discrete G_n-spectrum $(F_n \wedge X)$. Thus, homotopy fixed points for E_n come from homotopy fixed points for F_n.

Theorem 9.7. *For any closed subgroup G of G_n and any spectrum X with trivial G-action, there is an isomorphism*

$$(\hat{L}(E_n \wedge X))^{hG} \cong \hat{L}((F_n \wedge X)^{hG})$$

in the stable category. In particular, $E_n^{hG} \cong \hat{L}(F_n^{hG})$.

Proof. After switching M_I and X, $(\hat{L}(E_n \wedge X))^{hG} \cong \operatorname{holim}_I (F_n \wedge X \wedge M_I)^{hG}$. By Remark 7.15, $(F_n \wedge X \wedge M_I)^{hG} \cong (F_n \wedge X)^{hG} \wedge M_I \simeq ((F_n \wedge X)^{hG} \wedge M_I)_{\mathtt{f}}$, where the isomorphism signifies a zigzag of natural weak equivalences. Thus,

$$(\hat{L}(E_n \wedge X))^{hG} \cong \operatorname{holim}_I ((F_n \wedge X)^{hG} \wedge M_I)_{\mathtt{f}} \cong \hat{L}((F_n \wedge X)^{hG}),$$

since, by Lemma 9.6, $(F_n \wedge X)^{hG}$ is $E(n)$-local. □

Corollary 9.8. *If X is a finite spectrum of type n, then there is an isomorphism $(F_n \wedge X)^{hG} \cong E_n^{hG} \wedge X$. In particular, $(F_n \wedge M_I)^{hG} \cong E_n^{hG} \wedge M_I$.*

Proof. Since X is finite, by Remark 7.15, $(F_n \wedge X)^{hG} \cong F_n^{hG} \wedge X$. Then, since F_n^{hG} is $E(n)$-local by Lemma 9.6, there are isomorphisms $F_n^{hG} \wedge X \cong \hat{L}(F_n^{hG}) \wedge X \cong E_n^{hG} \wedge X$. □

We conclude this section by observing that smashing with a finite spectrum commutes with taking the homotopy fixed points of E_n.

Theorem 9.9. *Let G be a closed subgroup of G_n and let X be a finite spectrum. Then there is an isomorphism $(E_n \wedge X)^{hG} \cong E_n^{hG} \wedge X$.*

Proof. Recall that $(E_n \wedge X)^{hG} = \operatorname{holim}_I (F_n \wedge M_I \wedge X)^{hG}$. The zigzag of natural weak equivalences between $(F_n \wedge M_I \wedge X)^{hG}$ and $(F_n \wedge M_I)^{hG} \wedge X$ yields

$$(E_n \wedge X)^{hG} \cong \operatorname{holim}_I ((F_n \wedge M_I)^{hG} \wedge X)_{\mathtt{f}} \simeq (\operatorname{holim}_I (F_n \wedge M_I)^{hG}) \wedge X,$$

where the weak equivalence is due to Lemma 2.7. □

10. The descent spectral sequence for $(\hat{L}(E_n \wedge X))^{hG}$

By applying the preceding two sections, it is now an easy matter to build the descent spectral sequence for $(\hat{L}(E_n \wedge X))^{hG}$.

Definition 10.1. Let X be a spectrum. If the tower $\{\pi_t(E_n \wedge M_I \wedge X)\}_I$ of abelian groups satisfies the Mittag-Leffler condition for all $t \in \mathbb{Z}$, then X is an *E_n-Mittag-Leffler spectrum.* If X is an E_n-Mittag-Leffler spectrum, then, for convenience, we say that X is E_n-ML.

Any finite spectrum X is E_n-ML, since $\{\pi_t(E_n \wedge M_I \wedge X)\}_I$ is a tower of finite abelian groups, by Lemma 2.13. However, an E_n-ML spectrum need not be finite. For example, for $j \geq 1$, let $X = E_n^{(j)}$. Then $\pi_t(E_n \wedge M_I \wedge X) \cong \mathrm{Map}_c^{\ell}(G_n^j, \pi_t(E_n)/I)$. Since $\{\pi_t(E_n)/I\}$ is a tower of epimorphisms, the tower $\{\mathrm{Map}_c^{\ell}(G_n^j, \pi_t(E_n)/I)\}$ is also, and $E_n^{(j)}$ is E_n-ML.

Theorem 10.2. *Let G be a closed subgroup of G_n and let X be any spectrum with trivial G-action. Let*

$$E_2^{s,t} = \pi^s\pi_t(\operatorname{holim}_I (\Gamma_G^{\bullet}(F_n \wedge M_I \wedge X)_{f,G})^G).$$

Then there is a conditionally convergent descent spectral sequence

$$E_2^{s,t} \Rightarrow \pi_{t-s}((\hat{L}(E_n \wedge X))^{hG}). \tag{10.3}$$

If X is E_n-ML, then $E_2^{s,t} \cong H^s_{\mathrm{cts}}(G; \pi_t(\hat{L}(E_n \wedge X)))$. In particular, if X is a finite spectrum, then descent spectral sequence (10.3) has the form

$$H^s_c(G; \pi_t(E_n \wedge X)) \Rightarrow \pi_{t-s}((E_n \wedge X)^{hG}).$$

Proof. As in Remark 9.4, $\hat{L}(E_n \wedge X) \cong \operatorname{holim}_I (F_n \wedge M_I \wedge X)_{f,G}$ is a continuous G-spectrum. Then (10.3) follows from Theorem 8.10 by considering the tower of spectra $\{(F_n \wedge M_I \wedge X)_f\}_I$. When X is E_n-ML, $\{(F_n \wedge M_I \wedge X)_f\}$ is a Mittag-Leffler tower of spectra, and thus, the simplification of the E_2-term in this case follows from Theorem 8.10. By Corollary 2.16, when X is finite, there is an isomorphism $H^s_{\mathrm{cts}}(G; \pi_t(E_n \wedge X)) \cong H^s_c(G; \pi_t(E_n \wedge X))$. ☐

As discussed in Remark 8.12, Theorem 8.6 gives a spectral sequence with abutment $\pi_*((\hat{L}(E_n \wedge X))^{hG})$, the same as the abutment of (10.3), but with E_2-term $H^s_{\mathrm{cont}}(G; \{\pi_t(E_n \wedge M_I \wedge X)\})$, which is, in general, different from the E_2-term of (10.3). We are interested in the descent spectral sequence of Theorem 10.2, not just because it is a second descent spectral sequence with an interesting E_2-term, but, as mentioned in §8, it can be compared with the $K(n)$-local E_n-Adams spectral sequence. (This comparison is work in progress.)

We conclude this paper with a computation that uses spectral sequence (10.3) to compute $\pi_*((\hat{L}(E_n \wedge E_n^{(j)}))^{hG_n})$, where $j \geq 1$ and G_n acts only on the leftmost factor. By Theorem 2.4,

$$\begin{aligned}
\pi_t(\hat{L}(E_n^{(j+1)})) &\cong \mathrm{Map}_c^{\ell}(G_n^j, \pi_t(E_n)) \\
&\cong \lim_I \mathrm{Map}_c^{\ell}(G_n, \mathrm{Map}_c(G_n^{j-1}, \pi_t(E_n \wedge M_I))) \\
&\cong \lim_I \mathrm{Map}_c(G_n, \mathrm{Map}_c(G_n^{j-1}, \pi_t(E_n \wedge M_I))),
\end{aligned}$$

where the tower $\{\mathrm{Map}_c(G_n, \mathrm{Map}_c(G_n^{j-1}, \pi_t(E_n \wedge M_I)))\}$ satisfies the Mittag-Leffler condition, by Remark 2.11. Thus, in spectral sequence (10.3), Theorem 2.15 implies that

$$\begin{aligned} E_2^{s,t} &\cong H^s_{\mathrm{cts}}(G_n; \pi_t(\hat{L}(E_n^{(j+1)}))) \\ &\cong \lim{}_I\, H^s_c(G_n; \mathrm{Map}_c(G_n, \mathrm{Map}_c(G_n^{j-1}, \pi_t(E_n \wedge M_I)))), \end{aligned}$$

which vanishes for $s > 0$, and equals $\mathrm{Map}_c(G_n^{j-1}, \pi_t(E_n))$, when $s = 0$. Thus,

$$\pi_*((\hat{L}(E_n^{(j+1)}))^{hG_n}) \cong \mathrm{Map}_c(G_n^{j-1}, \pi_*(E_n)) \cong \pi_*(\hat{L}(E_n^{(j)})),$$

as abelian groups. Therefore, for $j \geq 1$, there is an isomorphism $(\hat{L}(E_n^{(j+1)}))^{hG_n} \cong \hat{L}(E_n^{(j)})$, in the stable category. The techniques described in this paper do not allow us to handle the $j = 0$ case, which would say that $E_n^{hG_n}$ and $\hat{L}(S^0) \simeq E_n^{\mathrm{h}G_n}$ are isomorphic.

References

[1] A. K. Bousfield. The localization of spectra with respect to homology. *Topology*, 18(4):257–281, 1979.

[2] A. K. Bousfield and E. M. Friedlander. Homotopy theory of Γ-spaces, spectra, and bisimplicial sets. In *Geometric applications of homotopy theory (Proc. Conf., Evanston, Ill., 1977), II*, volume 658 of *Lecture Notes in Math.*, pages 80–130. Springer, Berlin, 1978.

[3] A. K. Bousfield and D. M. Kan. *Homotopy limits, completions and localizations*. Springer-Verlag, Berlin, 1972. Lecture Notes in Mathematics, Vol. 304.

[4] Ethan S. Devinatz. Morava modules and Brown-Comenetz duality. *Amer. J. Math.*, 119(4):741–770, 1997.

[5] Ethan S. Devinatz. The generating hypothesis revisited. In *Stable and unstable homotopy (Toronto, ON, 1996)*, pages 73–92. Amer. Math. Soc., Providence, RI, 1998.

[6] Ethan S. Devinatz and Michael J. Hopkins. Homotopy fixed point spectra for closed subgroups of the Morava stabilizer groups. *Topology*, 43(1):1–47, 2004.

[7] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal. *Analytic pro-p groups*. Cambridge University Press, Cambridge, second edition, 1999.

[8] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory*. American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole.

[9] Thomas Geisser and Lars Hesselholt. Topological cyclic homology of schemes. In *Algebraic K-theory (Seattle, WA, 1997)*, volume 67 of *Proc. Sympos. Pure Math.*, pages 41–87. Amer. Math. Soc., Providence, RI, 1999.

[10] P. G. Goerss and J. F. Jardine. Localization theories for simplicial presheaves. *Canad. J. Math.*, 50(5):1048–1089, 1998.

[11] Paul G. Goerss. Homotopy fixed points for Galois groups. In *The Čech centennial (Boston, MA, 1993)*, pages 187–224. Amer. Math. Soc., Providence, RI, 1995.

[12] Paul G. Goerss and Michael J. Hopkins. André-Quillen (co)-homology for simplicial algebras over simplicial operads. In *Une dégustation topologique [Topological morsels]: homotopy theory in the Swiss Alps (Arolla, 1999)*, pages 41–85. Amer. Math. Soc., Providence, RI, 2000.

[13] Paul G. Goerss and Michael J. Hopkins. Moduli spaces of commutative ring spectra. In *Structured ring spectra*. Cambridge University Press, November, 2004.

[14] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*. Birkhäuser Verlag, Basel, 1999.

[15] Michael J. Hopkins, Mark Mahowald, and Hal Sadofsky. Constructions of elements in Picard groups. In *Topology and representation theory (Evanston, IL, 1992)*, pages 89–126. Amer. Math. Soc., Providence, RI, 1994.

[16] Mark Hovey. Bousfield localization functors and Hopkins' chromatic splitting conjecture. In *The Čech centennial (Boston, MA, 1993)*, pages 225–250. Amer. Math. Soc., Providence, RI, 1995.

[17] Mark Hovey. *Model categories*. American Mathematical Society, Providence, RI, 1999.

[18] Mark Hovey, Brooke Shipley, and Jeff Smith. Symmetric spectra. *J. Amer. Math. Soc.*, 13(1):149–208, 2000.

[19] Mark Hovey and Neil P. Strickland. Morava K-theories and localisation. *Mem. Amer. Math. Soc.*, 139(666):viii+100, 1999.

[20] Uwe Jannsen. Continuous étale cohomology. *Math. Ann.*, 280(2):207–245, 1988.

[21] J. F. Jardine. Simplicial presheaves. *J. Pure Appl. Algebra*, 47(1):35–87, 1987.

[22] J. F. Jardine. Stable homotopy theory of simplicial presheaves. *Canad. J. Math.*, 39(3):733–747, 1987.

[23] J. F. Jardine. *Generalized étale cohomology theories*. Birkhäuser Verlag, Basel, 1997.

[24] J. F. Jardine. Generalised sheaf cohomology theories. In *Axiomatic, enriched and motivic homotopy theory*, volume 131 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 29–68. Kluwer Acad. Publ., Dordrecht, 2004.

[25] Michel Lazard. Groupes analytiques p-adiques. *Inst. Hautes Études Sci. Publ. Math.*, (26):389–603, 1965.

[26] Saunders Mac Lane and Ieke Moerdijk. *Sheaves in geometry and logic*. Springer-Verlag, New York, 1994. A first introduction to topos theory, Corrected reprint of the 1992 edition.

[27] Mark Mahowald and Hal Sadofsky. v_n-telescopes and the Adams spectral sequence. *Duke Math. J.*, 78(1):101–129, 1995.

[28] M. A. Mandell, J. P. May, S. Schwede, and B. Shipley. Model categories of diagram spectra. *Proc. London Math. Soc. (3)*, 82(2):441–512, 2001.

[29] Stephen A. Mitchell. Hypercohomology spectra and Thomason's descent theorem. In *Algebraic K-theory (Toronto, ON, 1996)*, pages 221–277. Amer. Math. Soc., Providence, RI, 1997.

[30] Stephen A. Mitchell. K-theory hypercohomology spectra of number rings at the prime 2. In *Une dégustation topologique [Topological morsels]: homotopy theory in the Swiss Alps (Arolla, 1999)*, volume 265 of *Contemp. Math.*, pages 129–157. Amer. Math. Soc., Providence, RI, 2000.

[31] Stephen A. Mitchell. The algebraic K-theory spectrum of a 2-adic local field. *K-Theory*, 25(1):1–37, 2002.

[32] Jack Morava. Noetherian localisations of categories of cobordism comodules. *Ann. of Math. (2)*, 121(1):1–39, 1985.

[33] Douglas C. Ravenel. *Nilpotence and periodicity in stable homotopy theory*. Princeton University Press, Princeton, NJ, 1992. Appendix C by Jeff Smith.

[34] Charles Rezk. Notes on the Hopkins-Miller theorem. In *Homotopy theory via algebraic geometry and group representations (Evanston, IL, 1997)*, pages 313–366. Amer. Math. Soc., Providence, RI, 1998.

[35] Luis Ribes and Pavel Zalesskii. *Profinite groups*. Springer-Verlag, Berlin, 2000.

[36] Stefan Schwede. S-modules and symmetric spectra. *Math. Ann.*, 319(3):517–532, 2001.

[37] Jean-Pierre Serre. Groupes analytiques p-adiques. In *Séminaire Bourbaki, 1963/64, Fasc. 2, Exposé 270*, page 10. Secrétariat mathématique, Paris, 1964.

[38] N. P. Strickland. Gross-Hopkins duality. *Topology*, 39(5):1021–1033, 2000.

[39] Peter Symonds and Thomas Weigel. Cohomology of p-adic analytic groups. In *New horizons in pro-p groups*, pages 349–410. Birkhäuser Boston, Boston, MA, 2000.

[40] John Tate. Relations between K_2 and Galois cohomology. *Invent. Math.*, 36:257–274, 1976.

[41] R. W. Thomason. Algebraic K-theory and étale cohomology. *Ann. Sci. École Norm. Sup. (4)*, 18(3):437–552, 1985.

[42] R. W. Thomason. The finite stable homotopy type of some topoi. *J. Pure Appl. Algebra*, 47(1):89–104, 1987.

[43] Stavros Tsalidis. On the étale descent problem for topological cyclic homology and algebraic K-theory. *K-Theory*, 21(2):151–199, 2000.

[44] Charles A. Weibel. *An introduction to homological algebra*. Cambridge University Press, Cambridge, 1994.

[45] John S. Wilson. *Profinite groups*. The Clarendon Press Oxford University Press, New York, 1998.